

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



18

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07D 263/24, 263/20, 413/10, 487/04, 413/04		A1	(11) International Publication Number: WO 97/37980
		(43) International Publication Date: 16 October 1997 (16.10.97)	
(21) International Application Number: PCT/US97/03458		(74) Agent: STEIN, Bruce; Pharmacia & Upjohn Company, Intellectual Property Legal Services, 301 Henrietta Street, Kalamazoo, MI 49001 (US).	
(22) International Filing Date: 28 March 1997 (28.03.97)			
(30) Priority Data: 60/015,499 11 April 1996 (11.04.96) US		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): PEARLMAN, Bruce, A. [US/US]; 3411 Willow Lake Drive #308, Kalamazoo, MI 49008 (US). PERRAULT, William, R. [US/US]; 5108 Pavilion Drive, Kalamazoo, MI 49001 (US). BARBACHYN, Michael, R. [US/US]; 1216 Miles Avenue, Kalamazoo, MI 49001 (US). MANNINEN, Peter, R. [US/US]; 1328 Edington Avenue, Portage, MI 49002 (US). TOOPS, Dana, S. [US/US]; 7123 Balfour Drive, Kalamazoo, MI 49002 (US). HOUSER, David, J. [US/US]; 10121 Lloy Street, Portage, MI 49002 (US). FLECK, Thomas, J. [US/US]; 6171 South 29th Street, Scotts, MI 49088 (US).		Published <i>With international search report.</i>	

(54) Title: **PROCESS TO PREPARE OXAZOLIDINONES**

(57) Abstract

The present invention includes processes for producing 5-hydroxymethyl substituted oxazolidinone alcohols (III) from carbamates (IIA) or a trifluoroacetamide (IIB) using a dihydroxy compound (I) or glycidol (IV) starting material and for the transformation of the hydroxymethyl-substituted oxazolidinone alcohols (III) to the corresponding amino compounds, 5-aminomethyl substituted oxazolidinone amines (VII) which are acylated to form commercially useful antibacterial 5-acylamidomethyl substituted oxazolidinone (VIII).

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROCESS TO PREPARE OXAZOLIDINONES

BACKGROUND OF THE INVENTION

5

1. Field of the Invention

The present invention relates to processes for producing 5-hydroxymethyl substituted oxazolidinone alcohols (III). Also disclosed is a process for transformation of the 5-hydroxymethyl substituted oxazolidinone alcohols (III) to the corresponding 5-aminomethyl substituted oxazolidinone amines (VII) which are useful in the production of oxazolidinone antibacterial pharmaceuticals (VIII).

10

2. Description of the Related Art

US Patents 5,164,510, 5,182,403 and 5,225,565 disclose 5-indolinyloxazolidinones, 3-(5'-indazolyl)oxazolidinones, 3-(fused-ring substituted)phenyloxazolidinones respectively useful as antibacterial agents.

15

US Patents 5,231,188 and 5,247,090 disclose various tricyclic [6.5.5] and [6.6.5]-fused ring oxazolidinones useful as antibacterial agents.

International Publication WO93/09103 discloses mono- and di-halo phenyl oxazolidinone anti-bacterials which are useful as pharmaceutical agents for their anti-bacterial action.

20

US 4,150,029, 4,250,318, 4,476,136, 4,340,606 and 4,461,773 disclose the synthesis of 5-hydroxymethyloxazolidinones from amines ($R-NHX_1$, where X_1 is -H or *p*-toluenesulfonyl) and R,S-glycidol ($C^*H_2-O-C^*H-CH_2-OH$ where the carbon atoms marked * are bonded together, cyclized to form an epoxide). The mixture of enantiomers produced by this process (represented by the formula $R-NH-CH_2-CHOH-CH_2-OH$) are separated by fractional crystallization of the mandelic acid salts. The enantiomerically pure R-diol is then converted into the corresponding 5R-hydroxymethyl substituted oxazolidinones (III) by condensation with diethylcarbonate in the presence of sodium methoxide. These 5R-hydroxymethyl substituted oxazolidinones are useful as synthetic precursors of pharmaceutically useful oxazolidinones. The large number of steps renders this process unattractive.

25

J. Med. Chem., 32, 1673 (1989), *Tetrahedron* 45, 1323 (1989) and US Patent 4,948,801 disclose a method of producing oxazolidinones which comprises reacting an isocyanate ($R-N=C=O$) with (R)-glycidyl butyrate in the presence of a catalytic amount of lithium bromide - tributylphosphine oxide complex to produce the corresponding 5R-butyryloxymethyl substituted oxazolidinone. The process is

30

- performed at 135-145°. The butyrate ester is then hydrolyzed in a subsequent step to give the corresponding 5-hydroxymethyl substituted oxazolidinone. The relative high cost and/or availability of the isocyanate starting material and requirement of high temperature detract significantly from the attractiveness of this method.
- 5 *Abstracts of Papers*, 206th National Meeting of the American Chemical Society, Chicago, IL, August, 1993; American Chemical Society: Washington, DC, 1993; ORGN 089; *J. Med. Chem.* 39, 673 (1996); *J. Med. Chem.* 39, 680 (1996); International Publications WO93/09103, WO93/09103, WO95/07271 and WO93/23384; PCT applications PCT/US95/12751 and PCT/US95/10992; *Abstracts of*
- 10 *Papers*, 35th Interscience Conference on Antimicrobial Agents and Chemotherapy, San Francisco, CA, September, 1995; American Society for Microbiology: Washington, DC, 1995; Abstract No. F208; *Abstracts of Papers*, 35th Interscience Conference on Antimicrobial Agents and Chemotherapy, San Francisco, CA, September, 1995; American Society for Microbiology: Washington, DC, 1995;
- 15 Abstract No. F207; *Abstracts of Papers*, 35th Interscience Conference on Antimicrobial Agents and Chemotherapy, San Francisco, CA, September, 1995; American Society for Microbiology: Washington, DC, 1995; Abstract No. F206; *Abstracts of Papers*, 35th Interscience Conference on Antimicrobial Agents and Chemotherapy, San Francisco, CA, September, 1995; American Society for
- 20 Microbiology: Washington, DC, 1995; Abstract No. F227; disclose the reaction of a carbamate with *n*-butyllithium, lithium diisopropylamide or lithium hexamethyldisilazide at -78° to -40° followed by glycidyl butyrate at -78° followed by warming to 20-25° to produce 5-hydroxymethyl substituted oxazolidinones (III) where the ester is cleaved during the reaction.
- 25 US Patents 4,062,862 and 4,236,012 disclose a process to prepare oxazolidinones which comprises reacting an epoxide with a primary (lacking any substituent on the nitrogen atom) carbamate in the presence of a catalyst. The process "is preferably conducted at a temperature of from 100° to 150° for several hours."
- 30 Canadian Patent 681,830 discloses a process to prepare oxazolidinones which comprises reacting an aryl ether of glycidol with a primary carbamate in the presence of an alkaline catalyst (preferably lithium amide or lithium hydroxide). The process was performed in the "preferred temperature range of 150° to 165°". The products are aryl ethers of 5-hydroxymethyl substituted oxazolidinones and the
- 35 yields are poor (40-78%).

J. Am. Chem. Soc., 64, 1291 (1942) and US Patent 3,547,951 disclose a method for converting primary alcohols to amines that involves treatment with methane sulfonyl chloride to produce the mesylate followed by contacting the mesylate with anhydrous ammonia at ambient temperature in a sealed reaction vessel under high pressure.

It is also known that the mesylates of primary alcohols react with aqueous ammonia to give the corresponding primary amines, but high temperature and high pressure (85 psig) are required. Normally this process cannot be used in ordinary general purpose reactors and must be run in special reactors rated for high pressure.

International Publication WO95/07271 discloses the ammonolysis of oxazolidinone mesylates.


US Patent 4,476,136 discloses a method of transforming 5-hydroxymethyl substituted oxazolidinones (III) to the corresponding 5(S)-aminomethyl substituted oxazolidinones (VII) that involves treatment with methane sulfonyl chloride followed by potassium phthalimide followed by hydrazine. This reaction sequence produces by-products which are difficult to separate from the desired product.

J. Med. Chem., 32, 1673 (1989) and *Tetrahedron* 45, 1323 (1989) disclose a method for transforming 5-hydroxymethylsubstituted oxazolidinones into the corresponding 5S-acetamidomethyl substituted oxazolidinones that involves treatment with methanesulfonyl chloride or tosyl chloride, followed by sodium azide, followed by trimethylphosphite or platinum dioxide/hydrogen, followed by acetic anhydride or acetyl chloride to give the desired 5(S)-acetamidomethyl substituted oxazolidinone. It is known that sodium azide is an explosion hazard.

US Patent 5,210,303 discloses the conversion of various substituted benzyl chlorides into the corresponding benzylamines by heating with aqueous ammonia in the presence of aromatic aldehydes to suppress dialkylation. The dialkylated impurity is generally difficult to remove, see *Chem. Lett.*, 1057 (1978).


SUMMARY OF INVENTION

Disclosed is a process to prepare 5-hydroxymethyl substituted oxazolidinones of formula (III)



where R_1 is


5



where X^1 is -H or -F;


where X^2 is -H or -F;

10 where Q^1 is:




15

b)




20

c)



25


d)



30


5

f)




10

g)




15

h)




20

i)




25

j)




30

k)




35

m)



Q^1 and X^2 taken together are:



5

where Z^1 is:

- a) $-CH_2-$,
- b) $-CH(R^4)-CH_2-$,
- c) $-C(O)-$, or
- 10 d) $-CH_2CH_2CH_2-$;

where Z^2 is:

- a) $-O_2S-$,
- b) $-O-$,
- c) $-N(R^7)-$,
- 15 d) $-OS-$, or
- e) $-S-$;

where Z^3 is:

- a) $-O_2S-$,
- b) $-O-$,
- 20 c) $-OS-$, or
- d) $-S-$;


where A^1 is:

- a) H- or
- b) CH_3- ;



25 where A^2 is:

- a) H-,
- b) $HO-$,
- c) CH_3- ,
- d) CH_3O- ,
- 30 e) $R^2O-CH_2-C(O)-NH-$,
- f) $R^3O-C(O)-NH-$,
- g) $(C_1-C_2)alkyl-O-C(O)-$,
- h) $HO-CH_2-$,
- i) CH_3O-NH- ,
- 35 j) $(C_1-C_3)alkyl-O_2C-$.

- k) $\text{CH}_3\text{-C(O)-}$,
 l) $\text{CH}_3\text{-C(O)-CH}_2\text{-}$,



10 A^1 and A^2 taken together are:



20 where R^1 is:

- a) -CHO ,
 b) -COCH_3 ,
 c) -COCHCl_2 ,
 25 d) -COCHF_2 ,
 e) $\text{-CO}_2\text{CH}_3$,
 f) $\text{-SO}_2\text{CH}_3$, or
 g) $\text{-COCH}_2\text{OH}$;

where R^2 is:

- 30 a) H- ,
 b) $\text{CH}_3\text{-}$,
 c) $\text{phenyl-CH}_2\text{-}$, or
 d) $\text{CH}_3\text{C(O)-}$;

where R^3 is:

- 35 a) $(\text{C}_1\text{-C}_3)\text{alkyl-}$, or

b) phenyl-;
where R^4 is:


- a) H-, or
b) HO-;

5 where R^5 is:

- a) H-,
b) (C_1-C_3) alkyl-,
c) $CH_2 = CH-CH_2-$ or
d) $CH_3-O-(CH_2)_2-$;


10 where R^6 is:

- a) $CH_3-C(O)-$,
b) $H-C(O)-$,
c) $Cl_2CH-C(O)-$,
d) $HOCH_2-C(O)-$,
15 e) CH_3SO_2- ,




- g) $F_2CHC(O)-$,

20



- i) $H_3C-C(O)-O-CH_2-C(O)-$,
j) $H-C(O)-O-CH_2-C(O)-$,

25




- l) $HC \equiv CH-CH_2O-CH_2-C(O)-$ or
30 m) phenyl- $CH_2-O-CH_2-C(O)-$;

where R^7 is:


- a) $R^2O-C(R^{10})(R^{11})-C(O)-$,
b) $R^3O-C(O)-$,
c) $R^8-C(O)-$,

35


5




10




15




20



25

where R^8 is:

30

35 where R^9 is:

- a) $-\text{CH}_3$,
 b) $-\text{CH}_2\text{Cl}$,
 c) $-\text{CH}_2\text{CH}=\text{CH}_2$,
 d) aryl or
 5 e) $-\text{CH}_2\text{CN}$;
 where R^{10} is H- or CH_3 -;
 where R^{11} is H- or CH_3 -;
 where R^{12} is:
 a) H-,
 10 b) $\text{CH}_3\text{O}-\text{CH}_2\text{O}-\text{CH}_2$ - or
 c) HOCH_2 -;
 where R^{13} is:
 a) CH_3 -,
 b) HOCH_2 -,
 15 c) $(\text{CH}_3)_2\text{N-phenyl}$, or
 d) $(\text{CH}_3)_2\text{N-CH}_2$ -;
 where R^{14} is:
 a) HO- ,
 b) $\text{CH}_3\text{O-}$,
 20 c) $\text{H}_2\text{N-}$,
 d) $\text{CH}_3\text{O-C(O)-O-}$,
 e) $\text{CH}_3\text{-C(O)-O-CH}_2\text{-C(O)-O-}$,
 f) $\text{phenyl-CH}_2\text{-O-CH}_2\text{-C(O)-O-}$,
 g) $\text{HO-(CH}_2)_2\text{-O-}$,
 25 h) $\text{CH}_3\text{O-CH}_2\text{-O-(CH}_2)_2\text{-O-}$, or
 i) $\text{CH}_3\text{O-CH}_2\text{-O-}$;
 where R^{15} is:
 a) H- or
 b) Cl- ;
 30 where R^{16} is:
 a) HO-
 b) $\text{CH}_3\text{O-}$, or
 c) F- ;
 where m is 0 or 1;
 35 where n is 1 thru 3;

where p is 0 or 1;

where aryl is phenyl substituted with zero (0) or one (1) of the following:

- a) -F,
- b) -Cl,
- 5 c) -OCH₃,
- d) -OH,
- e) -NH₂,
- f) -(C₁-C₄)alkyl,
- g) -O-C(O)-OCH₃, or

- 10 h) -NO₂ and protected forms thereof, which comprises contacting a hydroxy compound selected from the group consisting of:

(a) (S)-, (R)- dihydroxy compound of formula (I)



or any mixture thereof where M₁ is -Cl, -Br or -O-SO₂-φ-CH₃, or


- 15 (b) (S)-, (R)- glycidol (IV)



or any mixture thereof where the carbon atoms designated by an * are each bonded to the same oxygen atom (-O-) to form a three member ring, with a carbamate of formula (IIA)




or a trifluoroacetamide of formula (IIB)



in the presence of a lithium cation and a base whose conjugate acid has a pK_a of greater than about 8 where -O-M₂ is a base whose acid has a pK_a of between about 8 and about 24, and where R₁ is as defined above.

Also disclosed is a process to prepare 5-aminomethyl substituted oxazolidinone amines of formula (VII)

30




where R₁ is as defined above, which comprises:

- 35 (1) contacting 5-hydroxymethyl substituted oxazolidinone alcohols of formula

(III)



5



(III)

where R_1 is as defined above with a sulfonylating agent selected from the group consisting of compounds of formula (V_a-V_d)

10



where n_1 is 0, 1 or 2;

15

where n_2 is 0 thru 4 with the provisos that:

if n_1 is 0, n_2 is 2, 3 or 4,

if n_1 is 1, n_2 is 0 or 1,


if n_1 is 2, n_2 is 0;

where n_3 is 5 - ($n_1 + n_2$);

20


where M_3 is Cl- or Br- to produce the corresponding oxazolidinone sulfonate of formula (VI_a-VI_d)

25




(VIa or VIb)

30



(VIc)

35



(VIId)

and

(2) contacting the oxazolidinone sulfonate (VI_a-VI_d) with ammonia at a pressure of less than about 30 psig.

DETAILED DESCRIPTION OF THE INVENTION

5 The process to produce the 5-hydroxymethyl substituted oxazolidinone alcohols (III) can use either the non-cyclic (S)-, (R)- dihydroxy compounds of formula (I) or any mixture thereof or (S)-, (R)- glycidol (IV) or any mixture to couple with the carbamate (IIA) or a trifluoroacetamide of formula (IIB).

10 The 5-hydroxymethyl substituted oxazolidinone alcohols (III) are useful intermediates to produce 5-aminomethyl substituted oxazolidinone amines (VII) which can be acylated to prepare pharmaceutically useful 5-acylamidomethyl substituted oxazolidinone (VIII) antibacterial agents. Because of an enantiomeric center, 5(R)-, 5(S)-acylamidomethyl substituted oxazolidinones (VIII) and mixtures thereof can be produced. The 5-acylamidomethyl substituted oxazolidinone (VIII) (S)-enantiomer has antibacterial activity, the (R)-enantiomer does not. The 5(S)-aminomethyl substituted oxazolidinone amine (VII) enantiomer is produced from the 5(R)-hydroxymethyl substituted oxazolidinone alcohol (III) enantiomer which is produced from the (S)-dihydroxy compound (I) or (S)-glycidol (IV). Therefore, the desired and preferred enantiomeric sequence is to use enantiomerically pure (S)-dihydroxy compound (I) or (S)-glycidol (IV) to give (R)-5-hydroxymethyl substituted oxazolidinone alcohol (III) which is used to give enantiomerically pure (S)-5-aminomethyl substituted oxazolidinone amine (VII) which is transformed to enantiomerically pure (S)-5-acylamidomethyl substituted oxazolidinone (VIII). However, it is readily apparent to one skilled in the art that one could easily perform the identical process steps with the opposite enantiomeric forms and at any point in the process invert an undesired enantiomeric configuration to the desired one. Therefore, using the chemistry of the claimed process with any of the enantiomeric forms is considered equivalent to the claimed processes.

25 The dihydroxy compounds, M₁-CH₂-CH(OH)-CH₂-OH, of formula (I) and glycidol compounds, C^{*}H₂-C^{*}H-CH₂-OH, of formula (IV) where the carbon atoms designated by an * are each bonded to the same oxygen atom (-O-) to form a three member ring, are known to those skilled in the art or can be readily prepared from known compounds by methods known to those skilled in the art. It is preferred that the hydroxy starting material be the dihydroxy compound (I). It is preferred that 35 the dihydroxy compound (I) and the glycidol (IV) be the (S)-enantiomer. It is

preferred that M_1 is Cl-; it is preferred that the dihydroxy compound (I) be claim 5, which can be purchased commercially.

- The carbamates, R_1 -NH-CO-O- M_2 , of formula (IIA) and the trifluoroacetamide, R_1 -NH-CO-CF₃, of formula (IIB) are either known to those skilled in the art or can readily be prepared from known compounds by methods known to those skilled in the art. The nature of the leaving group M_2 is not important since it is lost during the course of the reaction as is known to those skilled in the art. Operable M_2 (leaving groups) are those where -O- M_2 is a base whose acid has a pK_a of between about 8 and about 24. Preferred M_2 includes
- 10 C_1 - C_{20} alkyl,
 - C_3 - C_7 cycloalkyl,
 - ϕ - optionally substituted with one or two C_1 - C_3 alkyl or F-, Cl-, Br-, I-,
 - $CH_2=CH-CH_2$ -,
 - $CH_3-CH=CH-CH_2$ -,
 - 15 $(CH_3)_2C=CH-CH_2$ -,
 - $CH_2=CH$ -,
 - $\phi-CH=CH-CH_2$ -,
 - $\phi-CH_2$ - optionally substituted on ϕ - with one or two -Cl, C_1 - C_4 alkyl, -NO₂,
 - CN, -CF₃,
 - 20 9-fluorenylmethyl,
 - $(Cl)_3C-CH_2$ -,
 - 2-trimethylsilylethyl,
 - $\phi-CH_2-CH_2$ -,
 - 1-adamantyl,
 - 25 $(\phi)_2CH$ -,
 - $CH\equiv C-C(CH_3)_2$ -,
 - 2-furanylmethyl,
 - isobornyl, more preferred leaving groups are C_1 - C_4 alkyl or benzyl. Any other leaving group which operates in a similar manner is considered equivalent to those identified above. The carbamate (IIA) and trifluoroacetamide (IIB) carry the aromatic/heteroaromatic group (R_1 -) of the 5-hydroxymethyl substituted oxazolidinone alcohol (III). It is preferred that R_1 is phenyl substituted with one -F and one substituted amino group; it is more preferred that R_1 is 3-fluoro-4-[4-(benzyloxycarbonyl)-1-piperazinyl]phenyl or 3-fluoro-4-(4-morpholinyl)phenyl.
 - 35 Depending on the particular substituents in R_1 , the groups may have to be protected

as is known to those skilled in the art, by means known to those skilled in the art to prevent undesirable side reactions. For example, if the R_1 substituent has a free primary or secondary hydroxy group, it is not necessary, but preferable to protect it with an alcohol protecting group in the formation of the 5-hydroxymethyl substituted oxazolidinone alcohols (III). However, an unprotected alcohol will in general interfere with the reaction of the dihydroxy compound (I) or glycidol (IV) with the carbamate (IIA) or trifluoroacetamide (IIB) to give the 5-hydroxymethyl substituted oxazolidinone alcohols (III). The unprotected alcohol will not in general interfere with the conversion of the 5-hydroxymethyl substituted oxazolidinone alcohols (III) to the corresponding 5-aminomethyl substituted oxazolidinone amines (VII), because it is very difficult or impossible to selectively protect a primary or secondary alcohol on the R_1 functionality in the presence of another primary or secondary alcohol. Suitable alcohol protecting groups are well known to those skilled in the art, preferred are C_1 - C_5 alkyl, ϕ - CH_2 -, CH_3 -O- CH_2 -, CH_3 -, CH_3 -S- CH_2 -, ϕ - CH_2 -O- CH_2 -, tetrahydropyranyl, CH_3CH (-O- C_2H_5)-, *p*-methoxybenzyl, *p*-methoxyphenyl, *p*-nitrobenzyl, $(\phi)_3C$ -, $(CH_3)_3Si$ -, $[CH_3-CH(CH_3)]_3Si$ -, $\phi(CH_3)_2Si$ -. These protecting groups are removed by means known to those skilled in the art. For example, if R_1 contains a hydroxy substituent, it must be protected during the transformation of the 5-hydroxymethyl substituted oxazolidinone alcohol (III) to the 5-aminomethyl substituted oxazolidinone amine (VII) or the 5-acylamidomethyl substituted oxazolidinone (VIII). If the R_1 substituent contains a free primary or secondary amino substituent it does not have to be protected during formation of the 5-hydroxymethyl substituted oxazolidinone alcohols (III) but must be protected during transformation of the 5-hydroxymethyl substituted oxazolidinone alcohols (III) to the corresponding 5-aminomethyl substituted oxazolidinone amines (VII) and the 5-acylamidomethyl substituted oxazolidinones (VIII). The reason is that the amino group will in general undergo an undesired side reaction during one or more of the steps involved in the transformation of the 5-hydroxymethyl substituted oxazolidinone alcohols (III) to the corresponding 5-acylamidomethyl substituted oxazolidinones (VIII). Therefore, it is preferable to protect any free amino substituent in the R_1 functionality prior to the reaction of the dihydroxy compound (I) or glycidol (IV) with the carbamate (IIA) or trifluoroacetamide (IIB). Amino protecting groups are very well known to those skilled in the art. Preferred amino protecting groups include:

(I) C_1 - C_4 alkyl,

(II) $\phi\text{-CH}_2\text{-}$,

(III) $(\phi)_3\text{C-}$,

(IV) $\text{R}_a\text{-CO-}$ where R_a is (A) H-, (B) $\text{C}_1\text{-C}_4$ alkyl, (C) $\text{C}_5\text{-C}_7$ cycloalkyl, (D) $(\text{C}_1\text{-C}_5 \text{ alkyl})\text{-O-}$, (E) $\text{Cl}_3\text{C-CH}_2\text{-O-}$, (F) $\text{H}_2\text{C=CH-CH}_2\text{-O-}$, (G) $\phi\text{-CH=CH-CH}_2\text{-O-}$, (H) $\phi\text{-CH}_2\text{-O-}$, (I) *p*-methoxyphenyl- $\text{CH}_2\text{-O-}$, (J) *p*-nitrophenyl- $\text{CH}_2\text{-O-}$, (K) $\phi\text{-O-}$, (L) $\text{CH}_3\text{-CO-CH}_2\text{-}$, (M) $(\text{CH}_3)_3\text{Si-O-}$,

(V) $\text{R}_b\text{-SO}_2\text{-}$ where R_b is: (A) $(\text{C}_1 \text{ alkyl})\text{-}$, (B) $\phi\text{-}$, (C) *p*-methylphenyl- and (D) $\phi\text{-CH}_2\text{-}$. A preferred amino protecting group is benzyloxycarbonyl which can be removed by catalytic hydrogenation as is known to those skilled in the art. There is nothing novel regarding the use of protecting groups in these reactions or the nature of the particular protecting groups. All this is well known to those skilled in the art. The protecting groups can be removed after the last reaction in which the protected substituent would be affected or carried along and removed after subsequent reactions as is known to those skilled in the art. For example, it may be preferable to carry the protecting group along until the final acylation step is completed, before removal, as is known to those skilled in the art. Optionally the R_1 substituent can be modified after the 5-acylamidomethyl substituted oxazolidinones (VIII) is produced depending on what chemical reactions are required as is known to those skilled in the art.

The reaction of either dihydroxy compounds (I) or glycidol (IV) with either the carbamates (IIA) or trifluoroacetamides (IIB) give the same 5-hydroxymethyl substituted oxazolidinone alcohols (III). The choice of whether to use a dihydroxy compound (I) or glycidol (IV) to produce a particular 5-hydroxymethyl substituted oxazolidinone alcohol (III) has to be made on a case by case basis. No starting material is preferred in all cases; there is no generally preferred way based on chemistry alone. The decision involves the commercial availability of the particular starting material, its chemical and enantiomeric purity, its cost, etc as is known to those skilled in the art.

One process of the present invention is the reaction of the dihydroxy compound (I) or glycidol (IV) with the carbamate (IIA) or trifluoroacetamides (IIB) in the presence of lithium cation (Li^+) and a base whose conjugate acid has a pK_a of greater than about 8.

The processes requires about one molar equivalent of either the dihydroxy compound (I) or glycidol (IV)/equivalent of carbamate (IIA) or trifluoroacetamides (IIB). The reaction requires a base, the nature of which is not critical so long as it is

strong enough to deprotonate the carbamate (II). Operable bases are those whose conjugate acid has a pK_a of greater than about 8. Preferred bases include compounds selected from the group consisting of:

- 5 alkoxy compounds of one thru seven carbon atoms,
- carbonate,
- methyl, *sec*-butyl and *t*-butyl carbanions,
- tri(alkyl)amines where the alkyl group is from 1 thru 4 carbon atoms,
- conjugate base of the carbamate (II),
- DBU,
- 10 DBN,
- N-methyl-piperidine,
- N-methyl morpholine,
- 2,2,2-trichloroethoxide and

- 15 $Cl_3C-CH_2-O^-$; most preferred bases are where the base is alkoxy of four or five carbon atoms. It is preferred that the four and five carbon alcohol bases be *t*-amylate or *t*-butoxide. Sodium or potassium bases in combination with a lithium salt (such as lithium chloride or lithium bromide) can be used forming the lithium cation and base *in situ*.

- 20 The nature of the solvent is not critical. Operable solvents include cyclic ethers such as THF, amides such as DMF and DMAC, amines such as triethylamine, acetonitrile, and alcohols such as *t*-amyl alcohol and *t*-butyl alcohol. The choice of solvent depends on the solubility of the carbamate (IIA) or trifluoroacetamide (IIB) as is known to those skilled in the art.

- 25 When the starting material is the dihydroxy compounds (I) it can be beneficial to react the dihydroxy compound (I) with an cyclizing agent prior to contacting with the carbamate (IIA) or trifluoroacetamide (IIB). The term "cyclizing agent" refers to a base that cyclizes the dihydroxy compound (I) to glycidol (IV). Operable cyclizing agents include bases whose acid has a pK_a of greater than about 7; preferred cyclizing agents are sodium, potassium or lithium butoxide, sodium or
- 30 potassium hydroxide, potassium carbonate, DBU, lithium, sodium and potassium amylate; most preferred is potassium *t*-butoxide. It is preferable to perform the reaction at $< 100^\circ$, more preferable to perform it at $< 70^\circ$, even more preferable to perform it at $< 50^\circ$ and most preferable to perform it at $< 25^\circ$. The reaction can be performed at room temperature (about 20 to about 25°). At about 20° , the reaction
- 35 requires about 8 hr to reach completion (in DMAC). If a faster reaction is desired,

the reaction can be run at higher temperature. As stated above, differentiation between primary alcohols and secondary alcohols is difficult. In the cyclization reaction, a simple alcohol is formed. For instance, benzyl alcohol is formed when a benzyl carbonate is subjected to the cyclization conditions. Removal of this alcohol is necessary to the success of the alcohol to amine conversion. This is accomplished by crystallization using ethyl acetate/heptane (1/2). The benzyl alcohol stays in solution and the desired oxazolidinone alcohol is isolated as a solid.

CHART C discloses the processes of transforming the 5-hydroxymethyl substituted oxazolidinone alcohols (III) to the corresponding 5-aminomethyl substituted oxazolidinone amines (VII). The situation of protecting the alcohol and/or amino groups on the R_1 functionality was discussed above. The 5-hydroxymethyl substituted oxazolidinone alcohols (III) are contacted with a sulfonylating agent (V_a - V_d) of four types. These are M_3 -SO₂-C₆H_{n3}(NO₂)_{n1}Cl_{n2} (V_a), O[-SO₂-C₆H_{n3}(NO₂)_{n1}Cl_{n2}]₂ (V_b), O(SO₂-F)₂ (V_c) and O(SO₂-CF₃)₂ (V_d). M_3 is a leaving group which includes Cl- or Br-; it is preferred that M_3 be Cl-. The 5-hydroxymethyl substituted oxazolidinones (III) are contacted with a sulfonylating agent (V_a - V_d) to form a oxazolidinone sulfonate (VI_a - VI_d) intermediate.

The sulfonation reaction of converting the 5-hydroxymethyl substituted oxazolidinones (III) to the corresponding oxazolidinone sulfonates (VI) is performed by contacting the 5-hydroxymethyl substituted oxazolidinones (III) with at least one molar equivalent of the sulfonylating agent (V_a - V_d) in the presence of a base in an inert solvent at about 0°. Operable bases include triethylamine, tributylamine, diisopropylethylamine, DABCO, DBU, DBN, *n*-butyl lithium, ethyl magnesium chloride and the equivalents thereof; preferred is triethylamine. Inert solvents include most organic solvents such as methylene chloride, THF, DMA, DMF, ethyl acetate, and the equivalent thereof; preferred is methylene chloride.

The ammonolysis reaction of the conversion of the oxazolidinone sulfonates (VI) to the corresponding 5-aminomethyl substituted oxazolidinone amines (VII) is performed under open or non-sealed conditions or under sealed conditions although it is preferred to be performed under sealed conditions.

In either case the ammonolysis reaction is carried out by contacting the oxazolidinone sulfonates (VI) with ammonia (preferably aqueous) preferably with a solvent or mixture of solvents. Preferred solvents are those that dissolve both the oxazolidinone sulfonates (VI) and the aqueous ammonia because by dissolving both contact between them is insured. However, the process is also operable with

solvents that only partially dissolve the oxazolidinone sulfonates (VI), the disadvantage is that the reaction in general is slower. In the case of the the *m*-nitrobenenesulfonates, the preferred solvent is a mixture of acetonitrile/isopropanol or THF/isopropanol. The system is put under reduced pressure. The system is then closed or sealed, and the ammonia (preferably aqueous ammonia) is added and heated to less than 50°, preferably to less than 40°, preferably to about 38° (about 3 psig). At about 38-40° the pressure is about 0 to about 10 psig, which is well below the ceiling pressure rating of general purpose reactors. Under these conditions, at about 60°, the psig is about 20. It is preferred that the ammonolysis reaction be performed at a pressure of about 0 to about 20 psig, preferably at about 0 to about 5 psig and at about 60° or less. Alternatively, the reaction is performed in an open system at reflux. In this case the temperature will be slightly lower and the reaction will require slightly longer to reach completion. The ammonia can be either aqueous, alcoholic or anhydrous; however, aqueous ammonia is preferred.

Alternatively, the contacting with aqueous ammonia can be performed in the presence of an aromatic aldehyde (IX, Ar-CHO), preferably salicylaldehyde. The 5-aminomethyl substituted oxazolidinone amines (VII) and the aldehyde (IX) form a Schiff base of the formula (oxazolidinone-N=CH-Ar) which is then hydrolyzed with aqueous acid, as is known to those skilled in the art, to give the desired 5-aminomethyl substituted oxazolidinone amines (VII). The aromatic aldehyde (IX) is useful in suppressing dimer formation.

The 5-aminomethyl substituted oxazolidinone amines (VII) are acylated by known means such as acyl halides or acyl anhydrides to form the corresponding 5-acylamidomethyl substituted oxazolidinone (VIII), see CHART D. Any alcohol or amino protecting groups must be removed after the 5-acylamidomethyl substituted oxazolidinones (VIII) are produced. However, they can be removed earlier in the reaction sequence depending on the particular substituents in question as is known to those skilled in the art.

The 5-acylamidomethyl substituted oxazolidinones (VIII) are known to be antibacterial pharmaceutical agents. R₂ is selected from the group consisting of -H, C₁-C₁₂ alkyl optionally substituted with one or more halogens, (C₃-C₇)cyclo(C₅-C₉)alkyl or -O-R_{2a} where R_{2a} is C₁-C₆ alkyl. It is preferred that R₂ is C₁ alkyl.

DEFINITIONS AND CONVENTIONS

The definitions and explanations below are for the terms as used throughout this entire document including both the specification and the claims.

I. CONVENTIONS FOR FORMULAS AND DEFINITIONS OF VARIABLES

The chemical formulas representing various compounds or molecular fragments in the specification and claims may contain variable substituents in addition to expressly defined structural features. These variable substituents are identified by a letter or a letter followed by a numerical subscript, for example, "Z_i" or "R_i" where "i" is an integer. These variable substituents are either monovalent or bivalent, that is, they represent a group attached to the formula by one or two chemical bonds. For example, a group Z₁ would represent a bivalent variable if attached to the formula CH₃-C(=Z₁)H. Groups R_i and R_j would represent monovalent variable substituents if attached to the formula CH₃-CH₂-C(R_i)(R_j)-H. When chemical formulas are drawn in a linear fashion, such as those above, variable substituents contained in parentheses are bonded to the atom immediately to the left of the variable substituent enclosed in parenthesis. When two or more consecutive variable substituents are enclosed in parentheses, each of the consecutive variable substituents is bonded to the immediately preceding atom to the left which is not enclosed in parentheses. Thus, in the formula above, both R_i and R_j are bonded to the preceding carbon atom. Also, for any molecule with an established system of carbon atom numbering, such as steroids, these carbon atoms are designated as C_i, where "i" is the integer corresponding to the carbon atom number. For example, C₆ represents the 6 position or carbon atom number in the steroid nucleus as traditionally designated by those skilled in the art of steroid chemistry. Likewise the term "R₆" represents a variable substituent (either monovalent or bivalent) at the C₆ position.

Chemical formulas or portions thereof drawn in a linear fashion represent atoms in a linear chain. The symbol "-" in general represents a bond between two atoms in the chain. Thus CH₃-O-CH₂-CH(R_i)-CH₃ represents a 2-substituted-1-methoxypropane compound. In a similar fashion, the symbol "=" represents a double bond, e.g., CH₂=C(R_i)-O-CH₃, and the symbol "≡" represents a triple bond, e.g., HC≡C-CH(R_i)-CH₂-CH₃. Carbonyl groups are represented in either one of two ways: -CO- or -C(=O)-, with the former being preferred for simplicity.

Chemical formulas of cyclic (ring) compounds or molecular fragments can be represented in a linear fashion. Thus, the compound 4-chloro-2-methylpyridine can be represented in linear fashion by N^{*}=C(CH₃)-CH=CCl-CH=C^{*}H with the convention that the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring. Likewise, the cyclic molecular fragment, 4-

(ethyl)-1-piperaziny] can be represented by $-N^*-(CH_2)_2-N(C_2H_5)-CH_2-C^*H_2$.

A rigid cyclic (ring) structure for any compounds herein defines an orientation with respect to the plane of the ring for substituents attached to each carbon atom of the rigid cyclic compound. For saturated compounds which have two substituents
 5 attached to a carbon atom which is part of a cyclic system, $-C(X_1)(X_2)-$ the two substituents may be in either an axial or equatorial position relative to the ring and may change between axial/equatorial. However, the position of the two substituents relative to the ring and each other remains fixed. While either substituent at times may lie in the plane of the ring (equatorial) rather than above or below the plane
 10 (axial), one substituent is always above the other. In chemical structural formulas depicting such compounds, a substituent (X_1) which is "below" another substituent (X_2) will be identified as being in the alpha (α) configuration and is identified by a broken, dashed or dotted line attachment to the carbon atom, i.e., by the symbol "- -" or "...". The corresponding substituent attached "above" (X_2) the other (X_1) is
 15 identified as being in the beta (β) configuration and is indicated by an unbroken line attachment to the carbon atom.

When a variable substituent is bivalent, the valences may be taken together or separately or both in the definition of the variable. For example, a variable R_i attached to a carbon atom as $-C(=R_i)-$ might be bivalent and be defined as oxo or
 20 keto (thus forming a carbonyl group $(-CO-)$ or as two separately attached monovalent variable substituents $\alpha-R_{i-j}$ and $\beta-R_{i-k}$. When a bivalent variable, R_i , is defined to consist of two monovalent variable substituents, the convention used to define the bivalent variable is of the form " $\alpha-R_{i-j}:\beta-R_{i-k}$ " or some variant thereof. In such a case both $\alpha-R_{i-j}$ and $\beta-R_{i-k}$ are attached to the carbon atom to give $-C(\alpha-R_{i-j})(\beta-R_{i-k})-$
 25 . For example, when the bivalent variable R_6 , $-C(=R_6)-$ is defined to consist of two monovalent variable substituents, the two monovalent variable substituents are $\alpha-R_{6-1}:\beta-R_{6-2}$, $\alpha-R_{6-9}:\beta-R_{6-10}$, etc, giving $-C(\alpha-R_{6-1})(\beta-R_{6-2})-$, $-C(\alpha-R_{6-9})(\beta-R_{6-10})-$, etc. Likewise, for the bivalent variable R_{11} , $-C(=R_{11})-$, two monovalent variable substituents are $\alpha-R_{11-1}:\beta-R_{11-2}$. For a ring substituent for which separate α and β
 30 orientations do not exist (e.g. due to the presence of a carbon carbon double bond in the ring), and for a substituent bonded to a carbon atom which is not part of a ring the above convention is still used, but the α and β designations are omitted.

Just as a bivalent variable may be defined as two separate monovalent variable substituents, two separate monovalent variable substituents may be defined
 35 to be taken together to form a bivalent variable. For example, in the formula

-C₁(R_i)H-C₂(R_j)H- (C₁ and C₂ define arbitrarily a first and second carbon atom, respectively) R_i and R_j may be defined to be taken together to form (1) a second bond between C₁ and C₂ or (2) a bivalent group such as oxa (-O-) and the formula thereby describes an epoxide. When R_i and R_j are taken together to form a more
5 complex entity, such as the group -X-Y-, then the orientation of the entity is such that C₁ in the above formula is bonded to X and C₂ is bonded to Y. Thus, by convention the designation "... R_i and R_j are taken together to form -CH₂-CH₂-O-CO- ..." means a lactone in which the carbonyl is bonded to C₂. However, when designated "... R_j and R_i are taken together to form -CO-O-CH₂-CH₂-the convention
10 means a lactone in which the carbonyl is bonded to C₁.

The carbon atom content of variable substituents is indicated in one of two ways. The first method uses a prefix to the entire name of the variable such as "C₁-C₄", where both "1" and "4" are integers representing the minimum and maximum number of carbon atoms in the variable. The prefix is separated from the variable
15 by a space. For example, "C₁-C₄ alkyl" represents alkyl of 1 through 4 carbon atoms, (including isomeric forms thereof unless an express indication to the contrary is given). Whenever this single prefix is given, the prefix indicates the entire carbon atom content of the variable being defined. Thus C₂-C₄ alkoxy carbonyl describes a group CH₃-(CH₂)_n-O-CO- where n is zero, one or two. By the second method the
20 carbon atom content of only each portion of the definition is indicated separately by enclosing the "C_i-C_j" designation in parentheses and placing it immediately (no intervening space) before the portion of the definition being defined. By this optional convention (C₁-C₃)alkoxy carbonyl has the same meaning as C₂-C₄ alkoxy carbonyl because the "C₁-C₃" refers only to the carbon atom content of the alkoxy group. Similarly while both C₂-C₆ alkoxy alkyl and (C₁-C₃)alkoxy(C₁-C₃)alkyl define
25 alkoxy alkyl groups containing from 2 to 6 carbon atoms, the two definitions differ since the former definition allows either the alkoxy or alkyl portion alone to contain 4 or 5 carbon atoms while the latter definition limits either of these groups to 3 carbon atoms.

30 When the claims contain a fairly complex (cyclic) substituent, at the end of the phrase naming/designating that particular substituent will be a notation in parentheses which will correspond to the same name/designation in one of the CHARTS which will also set forth the chemical structural formula of that particular substituent.

35

II. DEFINITIONS

All temperatures are in degrees Centigrade.

TLC refers to thin-layer chromatography.

THF refers to tetrahydrofuran.

5 DMF refers to dimethylformamide.

DBU refers to 1,8-diazabicyclo[5.4.0]undec-7-ene.

DBN refers to 1,5-diazabicyclo[4.3.0]non-5-ene.

DABCO refers to 1,4-diazabicyclo[2.2.2]octane.

DMA refers to dimethylacetamide.

10 Saline refers to an aqueous saturated sodium chloride solution.

Chromatography (column and flash chromatography) refers to purification/separation of compounds expressed as (support, eluent). It is understood that the appropriate fractions are pooled and concentrated to give the desired compound(s).

15 IR refers to infrared spectroscopy.

CMR refers to ^{13}C magnetic resonance spectroscopy, chemical shifts are reported in ppm (δ) downfield from TMS.

NMR refers to nuclear (proton) magnetic resonance spectroscopy, chemical shifts are reported in ppm (δ) downfield from tetramethylsilane.

20 $-\phi$ refers to phenyl (C_6H_5).

$[\alpha]_{\text{D}}^{25}$ refers to the angle of rotation of plane polarized light (specific optical rotation) at 25° with the sodium D line (589A).

MS refers to mass spectrometry expressed as m/e, m/z or mass/charge unit.

25 $[\text{M} + \text{H}]^+$ refers to the positive ion of a parent plus a hydrogen atom. EI refers to electron impact. CI refers to chemical ionization. FAB refers to fast atom bombardment.

HRMS refers to high resolution mass spectrometry.

30 Pharmaceutically acceptable refers to those properties and/or substances which are acceptable to the patient from a pharmacological/toxicological point of view and to the manufacturing pharmaceutical chemist from a physical/chemical point of view regarding composition, formulation, stability, patient acceptance and bioavailability.

When solvent pairs are used, the ratios of solvents used are volume/volume (v/v).

35 When the solubility of a solid in a solvent is used the ratio of the solid to the

solvent is weight/volume (wt/v).

NNNNNN-NN-N refers to Chemical Abstracts Service (CAS, Columbus, Ohio) registry numbers where each "N" is an integer from 0 thru 9, but deleting leading zeros in the 6-digit portion of the number. Registry numbers are assigned to a particular chemical compound by CAS criteria, provided that the compound has been found to exist and it has been characterized in some way. Compounds published from approximately 1967 to the present are registered publicly and the registry number is the key to finding references in the CAS data base for such a registered compound. The CAS data base is publicly available from several database vendors such as STN International, System Development Corporation (SDC) Orbit Search Service, Lockheed Dialog, Bibliographic Retrieval Systems, Questrel, etc. CAS registry numbers are included in the EXAMPLES for some of the compounds which have been registered.

"psig" refers to "gauge pressure" equal to pressure (in psi) minus 1 atmosphere (14.7 psi).

EXAMPLES

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, practice the present invention to its fullest extent. The following detailed examples describe how to prepare the various compounds and/or perform the various processes of the invention and are to be construed as merely illustrative, and not limitations of the preceding disclosure in any way whatsoever. Those skilled in the art will promptly recognize appropriate variations from the procedures both as to reactants and as to reaction conditions and techniques.

EXAMPLE 1 (R)-[N-3-[3-Fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol (III)

A mixture of N-carbobenzoxy-3-fluoro-4-(N-carbobenzoxypiperazinyl)aniline (II, *J. Med. Chem.*, 39(3), 673 (1996)), 100 g of 98.4% pure material, 0.2133 moles) in DMAC (300 ml) is cooled to 0°. In a separate flask, a mixture of *t*-amyl alcohol (75 ml, 60.37 g, 0.685 moles, 3.23 eq) and heptane (75 ml) is cooled to -10° and treated with *n*-butyllithium in heptane (290 ml, 203 g of 14.4% wt/v solution, containing 29.2 g or 0.456 moles = 2/15 eq of *n*-butyllithium), keeping the temperature below 10°. The lithium *t*-amylate mixture is then added to the N-carbobenzoxy-3-fluoro-4-(N-carbobenzoxypiperazinyl)aniline (II) keeping the temperature below 10°.

Neat S-(+)-3-chloro-1,2-propanediol (I, CAS #60827-45-4, 22 ml, 29.1 g, 0.263

moles, 1.24 eq) is then added, rinsing with a small amount of heptane. The reaction mixture is then stirred at 20-25° and monitored by TLC (methanol/methylene chloride; 5/95) until the reaction is complete. The reaction mixture is then added to a mixture of acetic acid (40 ml, 42.0 g, 0.699 moles, 3.29 eq) in methanol (700 ml) and water (700 ml). The slurry formed is stirred at 20-25° for 30 min, cooled to 0°, stirred at 0° for 30 min, and filtered. The cake is washed with methanol/water (50/50) and dried under reduced pressure to give the title compound, TLC (methylene chloride/methanol, 95/5) R_f = 0.43.

EXAMPLE 2 (R)-[N-3-[3-Fluoro-4-[N-1-(4-carbobenzoxypiperazinyl)phenyl]-2-oxo-5-oxazolidinyl]methanol (III)

t-Amyl alcohol (0.967g, 10.97mmol, 2.571 eq) is cooled to -10°. Butyl lithium (4.3 ml, 2.5 M in hexanes, 10.8 mmol, 2.5 eq) is added with agitation while maintaining the temperature at less than 5°.

N-Carbobenzoxo-3-fluoro-4-(N-carbobenzoxypiperazinyl)aniline (II, 1.9780 g, 4.267 mmol, 1.000 eq) and dimethylacetamide (6.2 ml) are mixed, agitated and cooled to -25° to give a thin slurry. The lithium *t*-amylate mixture is then added to the N-benzyloxycarbonyl-3-fluoro-4-((4-benzyloxycarbonyl)-1-piperazinyl)aniline (II) mixture while maintaining less than -20°. The resultant mixture is warmed to 0° and S-(+)-3-chloro-1,2-propanediol (I, 0.5672 g, 5.131 mmol, 1.20 eq) is added. The resultant mixture is warmed to 21° and stirred for 7.5 hrs. The reaction mixture is added to a methanol (28 ml) and glacial acetic acid (0.73 ml, 12.75 mmol) mixture at 20-22°. The resulting slurry is then cooled to -30° and the product collected by vacuum filtration and washed with -30° methanol. The solids are dried in a stream of nitrogen to give the title compound, TLC (eluant chloroform/methanol, 90/10), R_f = 0.67; CMR (CDCl₃) 43.91, 46.39, 50.58, 62.60, 67.29, 72.89, 107.21, 107.56, 113.85, 119.36, 127.92, 128.09, 128.52, 133.51, 133.65, 136.05, 136.17, 136.57, 153.91, 154.80, 155.25 and 157.17 δ ; NMR (CDCl₃) 7.43, 7.31-7.37, 7.09, 6.88, 5.15, 4.67-4.90, 3.89-3.99, 3.67-3.74, 3.66, 3.25 and 2.98 δ ; MS (CI, m/e) = 430 (100%, P+1).

EXAMPLE 3 (R)-[N-3-(3-Fluoro-4-(4-morpholinylphenyl)-2-oxo-5-oxazolidinyl]methanol (III)

Tetrahydrofuran (3.0 ml) and *t*-amyl alcohol (0.66 ml, 6.03 mmol, 2.00 eq) are mixed. Butyl lithium (1.8 ml, 2.5 M in hexanes, 4.55 mmol, 1.5 eq) is added with agitation and while maintaining less than 2.5°.

N-Carbobenzoxo-3-fluoro-4-morpholinylaniline (II, *J. Med. Chem.*, 39(3), 673

(1996), 0.9942 g, 3.009 mmol, 1.000 eq) and tetrahydrofuran (3.5 ml) are mixture agitated and cooled. The lithium *t*-amylate mixture is then added to the carbamate (II) mixture while maintaining the temperature less than 8° and rinsed in with tetrahydrofuran (1 ml).

- 5 Tetrahydrofuran (3.2 ml) and S-(+)-3-chloro-1,2-propanediol (I, 0.299 ml, 3.58 mmol, 1.19 eq) are mixed. The mixture is cooled to -16° and potassium *t*-butoxide (3.2 ml, 1.0 M in tetrahydrofuran, 3.2 mmol, 1.07 eq) is added while maintaining the temperature at less than -10°. The resulting slurry is stirred at -14 to 0° for 1 hr then added to the lithium anion mixture while maintaining both mixtures at 0°,
10 then rinsed in with THF (2 ml). The resultant slurry is stirred at 20-23° for 2 hr then cooled to 6° and a mixture of citric acid monohydrate (0.4459 g, 2.122 mmol, 0.705 eq) in water (10 ml) is added. The resultant liquid phases are separated and the lower aqueous phase is washed with ethyl acetate (12 ml). The organic layers are combined and solvent is removed under reduced pressure until a net weight of
15 9.73 g remains. Heptane (10 ml) and water (5 ml) are added and solvent is removed by reduced pressure until a total volume of 5 ml remains. The precipitated product is collected by vacuum filtration and washed with water (7 ml). The solids are dried in a stream of nitrogen to give the title compound, TLC (chloroform/methanol, 95/5) $R_f = 0.23$; CMR (CDCl_3) 46.42, 51.01, 62.58, 73.07, 107.29, 107.64, 113.94, 118.80,
20 118.85, 128.28, 128.61, 133.15, 133.29, 136.26, 136.38, 153.82, 154.92 and 157.08 δ ; NMR (CDCl_3) 7.42, 7.32-7.37, 7.10, 4.67-4.75, 3.90-4.00, 3.86, 3.70-3.73, 3.44 and 3.03 δ ; MS (EI, m/e) = 296.

Alternatively, the crude product can be extracted with methylene chloride. The solvent is removed under reduced pressure. The solids are redissolved in hot
25 ethyl acetate, heptane is added, the mixture is cooled and the title compound is recovered.

EXAMPLE 4 (R)-[N-3-[3-Fluoro-4-[N-1-(4-carbobenzoxypiperazinyl)phenyl]-2-oxo-5-oxazolidinyl]methanol (III)

- A solution of *t*-amyl alcohol (75 ml, 60.3 g, 0.68 m) and heptane (75 ml) is
30 stirred and cooled to -10°. The mixture is treated with *n*-butyl lithium in heptane (1.6 M, 0.46 m, 290 ml) over a 30 min period while maintaining a temp < 10°. After 30 min, the mixture of lithium *t*-amylate is added to a mixture of N-carbobenzoxo-3-fluoro-4-(N-carbobenzoxypiperazinyl)aniline (II, 100 g, 0.22 m) and dimethylacetamide (300 ml) at 0° while maintaining a temp < 10°. The mixture is
35 stirred 30 min, then treated with S-(+)-3-chloro-1,2-propanediol (I, 22 ml, 0.26 m).

The cooling is removed, and the mixture is allowed to warm to 20-25°. The reaction is monitored by TLC and is judged complete after about 8 hr. The reaction mixture is poured into a mixture of methanol (700 ml), water (700 ml) and acetic acid (40 ml) and stirred for 30 min at 20-25°, then stirred for 30 min with cooling to 0°. The mixture is filtered, washed with aqueous methanol (50/50) and dried under reduced pressure at 45° to give the title compound, TLC (silica gel; methanol/methylene chloride, 5/95) R_f = 0.5. (90.3% yield).

EXAMPLE 5 3-Nitrobenzenesulfonate ester (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]-phenyl]-2-oxo-5-oxazolidinyl]methanol (VI)

A mixture of (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol (III, EXAMPLE 1, 43 g, 0.1 m) and methylene chloride (500 ml) is treated with triethylamine (32 ml, 0.23 m) and cooled to -5°. To this mixture is added a mixture of 3-nitrobenzenesulfonyl chloride (CAS # 121-51-7, 32 g, 0.14 ml) in methylene chloride (60 ml) while maintaining the temp < 10° over a 1 hr period. The reaction is monitored by TLC and judged complete after 45 min. The mixture is diluted with methylene chloride (500 ml) and then washed with water (2 x 600 ml). The organic phase is then washed with hydrochloric acid (1N, 400 ml) and concentrated to a thick residue. The residue is diluted with methanol (200 ml) and stirred for 1.5 hr. The solids are filtered, washed with methanol and dried under reduced pressure at 40° overnight to give the title compound, TLC (silica gel; methanol/methylene chloride, 5/95) R_f = 0.75.

EXAMPLE 6 (S)-N-[[3-[3-Fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]-phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (VII)

A slurry of 3-nitrobenzenesulfonate ester (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]-phenyl]-2-oxo-5-oxazolidinyl]methanol (VI, EXAMPLE 5, 50 g, 0.081 ml), isopropanol (250 ml), acetonitrile (400 ml) and aqueous ammonium hydroxide (29% ammonia by wt, 500 ml) is heated at 40° for 3.5 hr. The mixture is then treated with more aqueous ammonia (100 ml) and stirred for 20 hr. The reaction is monitored by TLC and judged complete at this time. The mixture is concentrated under reduced pressure with heat and suspended in methylene chloride/water (1250 ml/750 ml). The phases are separated and the organic phase concentrated to give a residue.

The residue is dissolved in methylene chloride (2 l) and treated with triethylamine (20 ml, 0.14 m). The mixture is then treated with acetic anhydride

(10 ml, 0.11 m) at 20-25° over 10 min. The acetylation is monitored by TLC and judged complete after 15 min. The organic mixture is washed with water (2 x 400 ml) then concentrated to a solid. The solids are recrystallized from ethanol (400 ml), filtered and dried under reduced pressure to give the title compound, TLC (silica gel; methanol/methylene chloride, 5/95) R_f = 0.6.

EXAMPLE 7 (S)-N-[[3-[3-fluoro-4-(1-piperazinyl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide hydrochloride (intermediate)

A mixture of (S)-N-[[3-[3-fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (VII, EXAMPLE 6, 35 kg, 74.5 moles),
10 palladium on carbon (5%, 10 kg, 50% water wet), methanol (550 l) and tetrahydrofuran (250 l) is agitated at 22 to 42° under a 42-50 psi hydrogen atmosphere. After 31 hours TLC analysis indicated complete reaction and the hydrogen atmosphere was replaced with nitrogen. The catalyst is removed by filtration and the filtrate concentrated under vacuum to 100 l. To the resulting
15 mixture, cooled to 2°, is added methanol (50 l) then a mixture of methanol (100 l) and acetyl chloride (6.04 kg, 77 moles) at -2° to 6°. The resulting mixture is stirred 90 minutes then concentrated under vacuum to 60 l, diluted with acetone (100 l) and concentrated further to 100 l. The resulting slurry is diluted with acetone (200 l) and stirred 15 hr at 16°. The solids are collected on a filter, washed with acetone
20 (50 l) and dried under reduced pressure at 20-25° to give the desired product. It is dissolved in methanol (56 l) at 53°, diluted with acetone (150 l), stirred 30 minutes at 48° then cooled to 15° and stirred 18 hr. The solids are collected on a filter, washed with acetone (50 l) and dried under reduced pressure at 20-25° to give the title compound, NMR (CDCl₃) 7.56-7.45, 7.31, 7.12-6.86, 4.79, 4.09-4.0, 3.81, 3.62,
25 3.40-3.11 and 2.01 δ .

EXAMPLE 8 (S)-N-[[3-[3-fluoro-4-[4-(hydroxyacetyl)-1-piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]-acetamide sesquihydrate (VIII)

To a stirred mixture of (S)-N-[[3-[3-fluoro-4-(1-piperazinyl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide hydrochloride (EXAMPLE 7, 16.2 kg, 43.5 moles),
30 tetrahydrofuran (205 kg) and triethylamine (10.1 kg, 100 moles) is added acetoxyacetyl chloride (6.5 kg, 47.8 moles) in tetrahydrofuran (11.1 kg) over 35 minutes keeping the temperature at 22-23°. After 40 minutes, at which time TLC and HPLC analysis indicated complete formation of the acetoxyacetamide intermediate, the mixture is concentrated under reduced pressure to 30 l, diluted
35 with methanol (100 l) and concentrated to 30 l. To the residue is added methanol

(25 l) and an aqueous solution of potassium carbonate (5.6 kg in 56 l). The resulting mixture is stirred 20 hr at 22-25° at which time TLC and HPLC analysis indicates the reaction is complete. The pH is adjusted to 7-7.5 with hydrochloric acid (4 N, 14.3 l). The mixture is stirred 18 hr at 15-22° then 3 hrs at 2-5°. The solids are collected on a filter, washed with water (68 l) and dried at 20-25° with recycled nitrogen to give the desired product. The crude product is dissolved in water (225 l) at 60-70°, clarified through a 0.6 micron filter, diluted with water rinse (55 l) and stirred 17 hrs. at 15°. The solids are collected on a filter, washed with water at 15° and dried at 45° with recycled nitrogen to a water content of 0.33%. These solids are dissolved in a solution of ethyl acetate (143 l), methanol (65 l) and water (1.95 l) at 60-65°. The solution is cooled to 15-25° and stirred 16 hrs for crystallization. The solids are collected on a filter, washed with ethyl acetate (75 l) and dried with 45° nitrogen to give the desired product. The product is recrystallized two more times from water (147 l then 133 l) at 60-70°, clarified each time through a 0.6 micron filter and rinsed with water (40 l and 30 l). The solids are dried on the filter at 30° with recycled nitrogen to give, after deagglomeration through a mill, the title compound as the sesquihydrate (6.45% water), TLC (silica gel; methanol/methylene chloride, 5/95) $R_f = 0.45$; $[\alpha]_D = -20^\circ$ (c = 1.0, ethanol).

EXAMPLE 9 3-Nitrobenzenesulfonate ester (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]-phenyl]-2-oxo-5-oxazolidinyl]methanol (VI)

To a slurry of (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol (III, EXAMPLE 1, 5.086 g, 11.86 mmol) in methylene chloride (50 mL) and triethylamine (2.0 mL, 14.38 mmol) at 0° is added dropwise over 6 minutes a solution of 3-nitrobenzene sulfonyl chloride (V) in methylene chloride (0.356M, 33.4 mL, 11.89 mmol). After stirring for 3.25 hrs, an additional 3.4 mL (1.21 mmol) of the 0.356 M solution of 3-nitrobenzene sulfonyl chloride (V) is added. After stirring for 1.75 hrs, hydrochloric acid (1N, 50 mL) is added. The phases are separated and the aqueous phase is extracted with methylene chloride. The combined organic phases are washed with saline, dried over magnesium sulfate and concentrated. The concentrate is crystallized from hot methylene chloride/methanol to give the title compound, mp = 155-157°; NMR (CDCl₃, 400 MHz) 8.72, 8.51, 8.23, 7.81, 7.35, 7.01, 6.91, 5.17, 4.85, 4.44, 4.39, 4.09, 3.85, 3.68 and 3.01 δ ; CMR (CDCl₃, 100 MHz) 44.26, 46.81, 50.91, 67.64, 69.54, 69.91, 107.85, 114.32, 119.85, 123.55, 128.30, 128.47, 128.91, 129.15, 131.51, 133.71, 136.99,

137.70, 148.71, 153.62, 155.57 and 155.88 δ ; IR (mineral oil mull) 1744, 1703, 1528, 1520, 1367, 1347 and 1192 cm^{-1} ; MS (EI, M/Z) 614, 411, 107, 91, 79, 65 and 56; $[\alpha]_D = -78^\circ$ ($c = 0.9812$, CHCl_3); TLC (ethyl acetate/hexane, 3/1) $R_f = 0.43$.

EXAMPLE 10 2-nitrobenzenesulfonate ester (R)-[N-3-[3-Fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol (VI)

Following the general procedure of EXAMPLE 5 (for the 3-nitrobenzenesulfonyl ester, (VI)) and making non-critical variations, (R)-[N-3-[3-fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol (III, EXAMPLE 1, 1.106 g, 2.578 mmol) is treated with triethyl amine (0.54 mL, 3.882 mmol) and commercial grade 2-nitrobenzenesulfonyl chloride (V, 679 mg, 3.064 mmol) to give the title compound, NMR (CDCl_3 , 400 MHz) 8.15, 7.82, 7.37, 7.06, 6.94, 5.17, 4.89, 4.59, 4.50, 4.10, 3.98, 3.69 and 3.03 δ ; IR (mineral oil mull) 1757, 1697, 1517, 1445, 1423, 1376, 1237 and 1188 cm^{-1} ; MS (EI, M/Z; rel. abundance): 614 (18.3, M^+), 91 (100), 69 (23.8) and 56 (52.9); TLC (ethyl acetate/hexane, 3/1) $R_f = 0.31$.

EXAMPLE 11 2,4-dinitrobenzenesulfonate ester (R)-[N-3-[3-Fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol (VI)

Following the general procedure of EXAMPLE 5 (for the 3-nitrobenzenesulfonyl ester) and making non-critical variations (R)-[N-3-[3-Fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol (III, EXAMPLE 1, 1.094 g, 2.550 mmol) is treated with triethyl amine (0.55 mL, 3.950 mmol) and commercial grade 2,4-dinitrobenzenesulfonyl chloride (833 mg, 3.124 mmol) to give the title compound, NMR (CDCl_3 , 400 MHz) 8.59, 8.38, 7.35, 7.02, 5.17, 4.88, 4.74, 4.58, 4.10, 3.98, 3.71, and 3.05 δ ; IR (mineral oil mull) 1756, 1697, 1554, 1541, 1517, 1351, 1237 and 1189 cm^{-1} ; MS (FAB, M/Z, rel. abundance) 660 (21.3, $[\text{M}+\text{H}]^+$), 659 (24.2, M^+), 102 (76.5) and 91 (100); TLC (ethyl acetate/hexane, 3/1) $R_f = 0.41$.

EXAMPLE 12 (R)-[N-3-[3-Fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol 4-chlorobenzenesulfonate ester (VI)

To a slurry of (R)-[N-3-[3-fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol (III, EXAMPLE 1, 3.450 g, 8.034 mmol) in methylene chloride (40 ml) and triethylamine (2.55 ml, 18.3 mmol) at -12° is added 4-chlorobenzenesulfonyl chloride (V, Aldrich Chemical Co.- commercial, 2.298 g, 10.88

mmol) as a solid all at once. The mixture is stirred in a 0° bath for 2.5 hrs then washed with water (2 X 35 ml), and 1N hydrochloric acid (35 ml). The organic extracts are concentrated to 20 ml total volume and methanol (50 ml) is added. The precipitate is collected by vacuum filtration, washed with methanol, dried and
 5 redissolved in methylene chloride (55 ml). The mixture is concentrated to a slurry of 32 g weight and methanol (11 ml) is added. The precipitate is collected by vacuum filtration, washed with methanol and dried. The solids are then dissolved in methylene chloride (58 ml) and column chromatographed (silica column, 93 g 40-63 μ ; eluted with 450 ml each of following ethyl acetate/cyclohexane mixtures 25/75; 35/65; 45/55; 55/45; collect last 50% of eluent). The collected eluent is concentrated to 200 ml and 200 ml heptane is added. The precipitate is collected by vacuum
 10 filtration and dried to give the title compound; TLC (silica gel; methanol/ chloroform 5/95) R_f = 0.53; MS (FAB, M/Z) = 604.7 (100%, [P+H]⁺); NMR (DMSO-d₆, 300 MHz) 7.93, 6.7, 7.75, 7.48-7.32, 7.12-7.03, 5.12, 4.93-4.92, 4.40, 4.09, 3.69, 3.57 and 2.96 δ ; 15 CMR (DMSO-d₆, 75 MHz) 43.51, 45.84, 50.22, 66.33, 69.75, 70.75, 106.63, 114.08, 119.83, 127.59, 127.87, 128.43, 129.62, 130.00, 133.31, 133.63, 135.52, 136.84, 139.63, 153.54, 154.40 and 154.62 δ .

EXAMPLE 13 (R)-[N-3-[3-Fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol 2,5-dichlorobenzenesulfonate ester
 20 (VI)

To a slurry of (R)-[N-3-[3-fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol (III, EXAMPLE 1, 3.439 g, 8.008 mmol) in methylene chloride (40 ml) and triethylamine (2.55 ml, 18.3 mmol) at -8° is added 2,5-dichlorobenzenesulfonyl chloride (V, Aldrich Chemical Co. - commercial, 2.675 g,
 25 10.90 mmol) as a solid all at once. The mixture is stirred in a 0° bath for 2.5 hrs then washed with water (2 X 35 ml), and 1N hydrochloric acid (35 ml). The organic extracts are then concentrated to 12.0 g which is column chromatographed (silica column, 108 g, 40-63 μ ; eluted with 450 ml each of following ethyl acetate/cyclohexane mixtures 10/90, 20/80, 30/70, 40/60 and 60/40 collecting the last
 30 20% of eluent). The collected eluent is concentrated and 300 ml methanol is added. The precipitate is collected by vacuum filtration, washed with methanol and dried to give the title compound, TLC (silica gel; methanol/chloroform 5/95) R_f = 0.66; MS (FAB, M/Z) = 638.6 (100%, [P+H]⁺); NMR (CDCl₃, 300 MHz) 8.04, 7.57-7.32, 7.06; 6.91, 5.16, 4.89-4.47, 4.42, 4.08, 3.93, 3.67 and 3.01 δ ; CMR (CDCl₃, 75 MHz) 43.93,
 35 45.51, 50.56, 67.26, 69.16, 69.46, 107.55, 113.98, 119.41, 127.92, 128.10, 128.54,

131.21, 131.46, 132.97, 133.44, 133.50, 134.68, 135.15, 136.45, 136.61, 153.36, 155.22 and 155.53 δ .

EXAMPLE 14 (R)-[N-3-[3-Fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol 4-nitrobenzenesulfonate ester (VI)

5 To a slurry of (R)-[N-3-[3-fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol (III, EXAMPLE 1, 3.437 g, 8.003 mmol) and 4-nitrobenzenesulfonyl chloride (V, 75% pure technical material, Aldrich Chemical Co.-commercial, 3.077 g, 10.41 mmol) in methylene chloride (32 ml) at 0° is added triethylamine (2.23 ml, 16.0 mmol). The mixture is stirred in a 0° bath for 1 hr then
10 water (1 ml) is added and the mixture stirred at 20-25° for 30 min. Methylene chloride (75 ml) is added and the mixture washed with hydrochloric acid (5%, 50 ml), then sodium bicarbonate (5%, 50 ml) and dried on magnesium sulfate. The organic extracts are then concentrated and the concentrate is taken up in boiling ethyl acetate/cyclohexane (1/1, 10 ml) and column chromatographed (silica gel, 4 cm
15 X 6", 40-63 μ ; eluting with about 400 ml each of following ethyl acetate/cyclohexane mixtures 20/80, 30/70, 40/60, 50/50, 60/40 and 70:30 collecting the last approximate 45% of eluent). The appropriate fractions are combined and concentrated to a solid which is dissolved in 70 ml methylene chloride and 50 ml ethyl acetate. The mixture is concentrated to 50 ml twice and cyclohexane (50 ml) is added after each
20 concentration. The precipitate is collected by vacuum filtration, washed with cyclohexane and dried to give the title compound, TLC (silica gel; ethyl acetate/cyclohexane 60/40) R_f = 0.37; NMR ($CDCl_3$, 300 MHz) 8.36, 8.07, 7.38-7.29, 7.03, 6.89, 5.15, 4.86-4.80, 4.39, 4.07, 3.80, 3.67 and 3.00 δ ; CMR ($CDCl_3$, 75 MHz) 43.85, 46.34, 50.45, 67.20, 69.17, 69.57, 107.64, 113.88, 119.34, 124.63, 127.85,
25 128.05, 128.49, 129.26, 132.67, 136.48, 136.57, 140.75, 150.95, 153.29, 155.14 and 155.40 δ .

EXAMPLE 15 (S)-[N-3-[3-Fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methylamine (VII)

Under nitrogen at 40° 3-nitrobenzenesulfonate ester (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]-phenyl]-2-oxo-5-oxazolidinyl]methanol (VI, EXAMPLE
30 5, 1.0099 g, 1.643 mmol), isopropanol (5.6 ml), acetonitrile (9.0 ml), benzaldehyde (0.50 ml, 4.92 mmol) and aqueous ammonia (29.8 wt% 9.5 ml, 148.6 mmol) are mixed. The mixture is stirred at 40° for 21.5 hrs, then concentrated under reduced pressure. Toluene (13.3 ml) and ethanol (6.0 ml) are added and the mixture warmed
35 in a 70° bath. Citric acid monohydrate (2.433 g, 11.58 mmol) was then added over

3.5 hrs and the phases separated at 64°. The organic phase is washed with water (2.5 ml) at 64°. The combined aqueous layers are washed with toluene (10 ml) at 64°. Toluene (10 ml) is then added to the aqueous and the mixture cooled to 0°. The precipitate is collected by vacuum filtration, washed with 0° toluene (10 ml) and 0° water (10 ml) and dried to a solid. A portion of this solid (0.7301 g) is slurried in water (10 ml) and methylene chloride (10 ml) and the pH adjusted from 2.78 to 13.92 with aqueous sodium hydroxide (50%, 0.3915 g, 4.90 mmol) at -4 to -2°. The mixture is warmed to 20-25° and sonicated with stirring for 0.5 hr. Methylene chloride (55 ml), saturated aqueous sodium chloride (5 ml) and water (35 ml) is added and the phases separated. The aqueous phase is washed two times with methylene chloride (25 ml) and the combined organics dried on sodium sulfate, filtered and concentrated under reduced pressure. Toluene (5 ml) is added followed by a slow addition of heptane (25 ml). The resultant precipitate is collected by vacuum filtration, washed with heptane (20 ml) and dried to give the title compound, TLC (silica gel; methanol/chloroform 10/90) $R_f = 0.32$; MS (EI), M/Z (relative intensity) = 428 (28%, M^+), 252 (15%), 92 (32%), 91 (100%); NMR ($CDCl_3$, 300 MHz) 7.46, 7.38-7.27, 7.12, 6.90, 5.16, 4.69-4.60, 3.98, 3.80, 3.67, 3.09, 3.00-2.92 and 1.30 δ ; CMR ($CDCl_3$, 75 MHz) 43.94, 44.89, 47.60, 50.63, 67.23, 73.84, 107.29, 113.72, 119.37, 127.92, 128.07, 128.52, 133.79, 136.05, 136.64, 154.57, 155.19 and 155.61 δ .

EXAMPLE 16 (R)-[N-3-[3-Fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 4-nitrobenzenesulfonate ester (VI)

To a slurry of (R)-[N-3-[3-fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol (III, EXAMPLE 3, 43.0 g, 145 mmol) and triethylamine (36 g, 355 mmol) in methylene chloride (450 ml) at 0° is added a mixture of 4-nitrobenzenesulfonyl chloride (V, 32 g, 145 mmol) in methylene chloride (55 ml). The mixture is stirred in a 0° bath for 30 min and then quenched with hydrochloric acid (10%, 200 ml). The organic phase is separated, and the aqueous phase is extracted again with methylene chloride (200 ml). The combined organic extracts are then concentrated column chromatographed (silica gel, 4 cm X 6", 40-63 μ ; methanol/methylene chloride 1-2/98-99, about 8 l). The appropriate fractions are combined and concentrated to give the title compound, $R_f = 0.2$; NMR ($CDCl_3$, 300 MHz) 8.73, 8.54, 8.23, 7.82, 7.33, 7.04, 6.91, 4.86, 4.42, 4.12, 3.86, and 3.05 δ ; CMR ($CDCl_3$, 75 MHz, partial) 46.42, 50.89, 66.87, 69.09, 69.45, 107.45, 113.95, 118.84, 123.14, 128.73, 131.08, 133.28 and 137.27 δ .

EXAMPLE 17 (S)-[N-3-[3-Fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methylaniline salicylaldehyde imine

A mixture of (R)-[N-3-(3-fluoro-4-(4-morpholinylphenyl)-2-oxo-5-oxazolidinyl]methanol 3-nitrobenzenesulfonate ester (VI, EXAMPLE 16, 20.608 g, 42.803 mmol), isopropanol (149 ml), acetonitrile (245 ml), salicylaldehyde (13.7 ml, 129 mmol) and aqueous ammonia (30%, 257 ml, 4.02 mol), is heated to 40° and stirred at 39-42° for 24 hrs. The mixture is then cooled to -22° and the precipitate collected by vacuum filtration, washed with water (10 ml) and dried to give the title compound, TLC (silica gel; methanol/chloroform 5/95) R_f = 0.79; EIMS (m/z, relative intensity) = 399 (M^+ , 51) 234 (11), 196 (11), 149 (22), 135 (100), 134 (47); NMR (300 MHz, $CDCl_3$) 8.44, 7.41, 7.33-6.87, 4.96-4.88, 4.12, 3.94-3.84 and 3.04 δ ; CMR ($CDCl_3$, 75 MHz) 48.21, 50.99, 61.94, 66.95, 71.30, 107.68, 114.12, 117.02, 118.43, 118.82, 119.01, 131.93, 133.04, 136.51, 154.24, 155.47, 160.78 and 168.87 δ .

EXAMPLE 18 (S)-N-[[3-(3-Fluoro-4-morpholinylphenyl)-2-oxo-5-oxazolidinyl]methyl]acetamide (VIII)

(S)-[N-3-[3-Fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methylaniline salicylaldehyde imine (EXAMPLE 17, 1.0068 g, 2.521 mmol) is slurried in water (10 ml) and 37% aqueous hydrochloric acid (0.417 ml, 5.04 mmol) and stirred at 20-25° for 15 hrs. Toluene (10 ml) is added and the phases separated; then, the organic phase is washed with hydrochloric acid (1M, 5 ml) and the combined aqueous phases are washed with toluene (10 ml). The toluene wash is back-extracted with hydrochloric acid (1M, 5 ml). The combined aqueous phases are then adjusted to pH 13.0 with aqueous sodium hydroxide (50%, 1.83 g, 22.9 mmol). To the resultant slurry is then added methylene chloride (10 ml) and sodium chloride (1 g) and the phases separated. The aqueous phase is then washed with methylene chloride (10 ml). To the combined organic phases is then added acetic anhydride (0.472 ml, 5.00 mmol) while maintaining 24-27°. The mixture is stirred 40 min, then water is added (5 ml). The phases are separated and the aqueous phase is washed with methylene chloride (5 ml). The combined organic phases are concentrated and ethyl acetate (25 ml) is added. The mixture is warmed to 70° and then the resultant mixture is slowly cooled to -25°. The precipitate is collected by vacuum filtration, washed with -25° ethyl acetate (5 ml) and dried to give the title compound, HPLC major component (99.93 area % at 254 nm detection) retention time = 0.97 min, column = Zorbax RX-C8, 250 X 4.6mm, mobile phase = 650 ml acetonitrile, 1.85 ml triethylamine, 1.30 ml acetic acid and sufficient water to make 1000 ml; flow rate =

3ml/min.

EXAMPLE 19 (R)-[N-3-[3-Fluoro-4-[N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol (III)

A mixture of N-carbobenzoxy-3-fluoro-4-(N-carbobenzoxypiperazinyl)aniline (II, 2.014 g, 4.345 mmol) and THF (10 ml) is cooled to -20°. In a separate flask, a solution of t-amyl alcohol (0.71 ml, 6.48 mmol) in THF (10 ml) at -33° is treated with n-butyllithium in heptane (13.65 wt%, 2.53 g, 5.38 mmol) while maintaining the mixture at less than -20°. The resultant lithium t-amylate solution is then added to the N-carbobenzoxy-3-fluoro-4-(N-carbobenzoxypiperazinyl)aniline mixture while maintaining less than -20° and rinsed in with THF (4 ml). To the resulting mixture at -28° is then added S-glycidol (IV, 0.3360 g, 4.536 mmol). The mixture is then stirred at -20° for 1.5 hrs, then at -16° for 17 hrs, at -11° for 4 hrs then at -1° for 2 hrs. HPLC assay then showed the major component to have a retention time consistent with the title compound (90.4 area % at 254 nm detection; retention time = 1.30 min; column = Zorbax RX-C8, 250 X 4.6 mm; mobile phase = 650 ml acetonitrile, 1.85 ml triethylamine, 1.30 ml acetic acid and add sufficient water to make 1000 ml; flow rate = 3 ml/min) as did TLC (silica gel; methanol/chloroform 10/90) R_f = 0.60.

EXAMPLE 20 (R)-[N-3-[3-Fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 4-nitrobenzenesulfonate ester (VI)

Following the general procedure of EXAMPLE 16 and making non-critical variations but starting with 4-nitrobenzenesulfonyl chloride, the title compound is obtained.

EXAMPLE 21 (R)-[N-3-[3-Fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 2-nitrobenzenesulfonate ester (VI)

Following the general procedure of EXAMPLE 16 and making non-critical variations but starting with 2-nitrobenzenesulfonyl chloride, the title compound is obtained.

EXAMPLE 22 (R)-[N-3-[3-Fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 2,4-dinitrobenzenesulfonate ester (VI)

Following the general procedure of EXAMPLE 16 and making non-critical variations but starting with 2,4-dinitrobenzenesulfonyl chloride, the title compound is obtained.

EXAMPLE 23 (R)-[N-3-[3-Fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 4-chlorobenzenesulfonate ester (VI)

Following the general procedure of EXAMPLE 16 and making non-critical variations but starting with 4-chlorobenzenesulfonyl chloride, the title compound is obtained.

EXAMPLE 24 (R)-[N-3-[3-Fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 2,5-dichlorobenzenesulfonate ester (VI)

5


Following the general procedure of EXAMPLE 16 and making non-critical variations but starting with 2,5-dichlorobenzenesulfonyl chloride, the title compound is obtained.

10

CHART A

(I)

+




(IIA)

or



(IIB)



(III)

CHART B

where the carbon atoms designated by an * are each bonded to the same oxygen atom (-O-) to form a three member ring or epoxide







CHART C

5




(III)

+




(Va)




(Vb)

10




(Vc)




(Vd)

15




(VIa or b)

20




(VIc)

25



(VIId)

30




(VII)

35

CHART D

5




(VII)

10



15

20



(VIII)


25

30

CLAIMS

1. A process to prepare 5-hydroxymethyl substituted oxazolidinones of formula (III)


5



(III)

where R_1 is

10



15


where X^1 is -H or -F;

where X^2 is -H or -F;

where Q^1 is:


20

a)




25

b)




30

c)




35

d)




5

f)




10

g)




15

h)




20

i)



25


j)



30




5



10

Q¹ and X² taken together are:



15

where Z¹ is:

- 20 a) -CH₂-,
 b) -CH(R⁴)-CH₂-,
 c) -C(O)-, or
 d) -CH₂CH₂CH₂-;

where Z² is:

- 25 a) -O₂S-,
 b) -O-,
 c) -N(R⁷)-,
 d) -OS-, or
 e) -S-;

where Z³ is:

- 30 a) -O₂S-,
 b) -O-,
 c) -OS-, or
 d) -S-;


where A¹ is:

- 35 a) H- or

b) CH_3 ;

where A^2 is:


- 5 a) H-,
 b) HO-,
 c) CH_3 -,
 d) CH_3O -,
 e) $\text{R}^2\text{O}-\text{CH}_2-\text{C}(\text{O})-\text{NH}$ -,
 f) $\text{R}^3\text{O}-\text{C}(\text{O})-\text{NH}$ -,
 g) $(\text{C}_1-\text{C}_2)\text{alkyl}-\text{O}-\text{C}(\text{O})$ -,
 10 h) $\text{HO}-\text{CH}_2$ -,
 i) $\text{CH}_3\text{O}-\text{NH}$ -,
 j) $(\text{C}_1-\text{C}_3)\text{alkyl}-\text{O}_2\text{C}$ -,
 k) $\text{CH}_3-\text{C}(\text{O})$ -,
 l) $\text{CH}_3-\text{C}(\text{O})-\text{CH}_2$ -,
 15



20



25 A^1 and A^2 taken together are:



30



35



where R^1 is:

- 5 a) -CHO,
 b) -COCH₃,
 c) -COCHCl₂,
 d) -COCHF₂,
 e) -CO₂CH₃,
 f) -SO₂CH₃, or
 10 g) -COCH₂OH;

where R^2 is:

- a) H-,
 b) CH₃-,
 c) phenyl-CH₂-, or
 15 d) CH₃C(O)-;

where R^3 is:

- a) (C₁-C₃)alkyl-, or
 b) phenyl-;

where R^4 is:


- 20 a) H-, or
 b) HO-;

where R^5 is:


- a) H-,
 b) (C₁-C₃)alkyl-,
 25 c) CH₂ = CH-CH₂- or
 d) CH₃-O-(CH₂)₂-;

where R^6 is:




- a) CH₃-C(O)-,
 b) H-C(O)-,
 30 c) Cl₂CH-C(O)-,
 d) HOCH₂-C(O)-,
 e) CH₃SO₂-;




5



10




15




20 where R^7 is:




25




30



35



g) R^9-SO_2- ,



- 5
- i) $HO-CH_2-C(O)-$,
 j) $R^{16}-(CH_2)_2-$,
 10 k) $R^{13}-C(O)-O-CH_2-C(O)-$,
 l) $(CH_3)_2N-CH_2-C(O)-NH-$,
 m) $NC-CH_2-$ or
 n) $F_2-CH-CH_2-$;

where R^8 is:

- 15 a) $H-$,
 b) $(C_1-C_4)alkyl$,
 c) $aryl-(CH_2)_p$,
 d) CH_2C- ,
 e) Cl_2HC- ,
 20 f) FH_2C- ,
 g) F_2HC- or
 h) $(C_3-C_6)cycloalkyl$;

where R^9 is:

- a) $-CH_3$,
 25 b) $-CH_2Cl$,
 c) $-CH_2CH=CH_2$,
 d) $aryl$ or
 e) $-CH_2CN$;

where R^{10} is $H-$ or CH_3- ;

30 where R^{11} is $H-$ or CH_3- ;

where R^{12} is:

- a) $H-$,
 b) $CH_3O-CH_2O-CH_2-$ or
 c) $HOCH_2-$;
 35 where R^{13} is:

- a) CH_3 -,
 b) HOCH_2 -,
 c) $(\text{CH}_3)_2\text{N-phenyl}$, or
 d) $(\text{CH}_3)_2\text{N-CH}_2$ -,
 5 where R^{14} is:
 a) HO- ,
 b) $\text{CH}_3\text{O-}$,
 c) $\text{H}_2\text{N-}$,
 d) $\text{CH}_3\text{O-C(O)-O-}$,
 10 e) $\text{CH}_3\text{-C(O)-O-CH}_2\text{-C(O)-O-}$,
 f) $\text{phenyl-CH}_2\text{-O-CH}_2\text{-C(O)-O-}$,
 g) $\text{HO-(CH}_2)_2\text{-O-}$,
 h) $\text{CH}_3\text{O-CH}_2\text{-O-(CH}_2)_2\text{-O-}$, or
 i) $\text{CH}_3\text{O-CH}_2\text{-O-}$;
 15 where R^{15} is:
 a) H- or
 b) Cl- ;
 where R^{16} is:
 a) HO-
 20 b) $\text{CH}_3\text{O-}$, or
 c) F- ;
 where m is 0 or 1;
 where n is 1 thru 3;
 where p is 0 or 1;
 25 where aryl is phenyl substituted with zero (0) or one (1) of the following:
 a) -F ,
 b) -Cl ,
 c) -OCH_3 ,
 d) -OH ,
 30 e) -NH_2 ,
 f) $\text{-(C}_1\text{-C}_4\text{)alkyl}$,
 g) -O-C(O)-OCH_3 , or
 h) -NO_2 and protected forms thereof,
 which comprises contacting a hydroxy compound selected from the group consisting
 35 of:

(a) (S)-, (R)- dihydroxy compound of formula (I)




(I)

or any mixture thereof where M_1 is -Cl, -Br or -O-SO₂- ϕ -CH₃, or

(b) (S)-, (R)- glycidol (IV)

5




(IV)

or any mixture thereof where the carbon atoms designated by an * are each bonded to the same oxygen atom (-O-) to form a three member ring, with a carbamate of formula (IIA)



(IIA)

10 or a trifluoroacetamide of formula (IIB)



(IIB)

in the presence of a lithium cation and a base whose conjugate acid has a pK_a of greater than about 8 where -O- M_2 is a base whose acid has a pK_a of between about 8 and about 24, and where R_1 is as defined above.

15

2. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 1 where the hydroxy compound is the dihydroxy compound of formula (I).

20 3. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 2 where the dihydroxy compound (I) is the (S)- enantiomer.

4. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 1 where M_1 is -Cl.

25 5. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 4 where the dihydroxy compound (I) is (S)-(+)-3-chloro-1,2-propanediol.

6. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 1 where the dihydroxy compound (I) is contacted with a cyclizing agent
30 prior to be contacted with the carbamate (IIA) or trifluoroacetamide (IIB).

7. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 6 where the cyclizing agent is a base whose acid has a pK_a of greater than about 7.

35

8. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 7 where the cyclizing agent is sodium, potassium or lithium butoxide, sodium or potassium hydroxide, potassium carbonate, DBU, lithium, sodium and potassium amylate.

5

9. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 1 where the hydroxy compound is the glycidol (IV).

10. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 9 where the glycidol (IV) is the (S)-enantiomer.

11. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 1 where M_2 is selected from the group consisting of:

- C_1 - C_{20} alkyl,
- 15 C_3 - C_7 cycloalkyl,
- ϕ - optionally substituted with one or two:
 - C_1 - C_3 alkyl,
 - F-, Cl-, Br-, I-,
 - $CH_2=CH-CH_2$ -,
 - 20 $CH_3-CH=CH-CH_2$ -,
 - $(CH_3)_2C=CH-CH_2$ -,
 - $CH_2=CH$ -,
 - $\phi-CH=CH-CH_2$ -,
 - $\phi-CH_2$ - optionally substituted on ϕ - with one or two -Cl, C_1 - C_4 alkyl, $-NO_2$,
 - 25 -CN, $-CF_3$,
 - 9-fluorenylmethyl,
 - $(Cl)_3C-CH_2$ -,
 - 2-trimethylsilylethyl,
 - $\phi-CH_2-CH_2$ -,
 - 30 1-adamantyl,
 - $(\phi)_2CH$ -,
 - $CH=C-C(CH_3)_2$ -,
 - 2-furanylmethyl,
 - isobornyl.

35

12. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 11 where M_2 is C_1 - C_4 alkyl or benzyl.
13. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 1 where R_1 is phenyl substituted with one -F and one substituted amino group.
14. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 13 where R_1 is:
3-fluoro-4-[4-(benzyloxycarbonyl)-1-piperazinyl]phenyl or
3-fluoro-4-(4-morpholinyl)phenyl.
15. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 1 where the base is selected from the group consisting of
alkoxy compounds of one thru seven carbon atoms,
carbonate,
methyl, sec-butyl and t-butyl carbanions,
tri(alkyl)amines where the alkyl group is from 1 thru 4 carbon atoms,
conjugate base of the carbamate (II),
DBU,
DBN,
N-methyl-piperidine,
N-methyl morpholine and
2,2,2-trichloroethoxide.
 $Cl_3C-CH_2-O^-$.
16. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 15 where the base is alkoxy of four or five carbon atoms.
17. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 1 where the protected form of R_1 is the alcohol protecting group selected from the group consisting of:
 C_1 - C_5 alkyl,
 $\phi-CH_2-$.

CH₃-O-CH₂-,
 CH₃-,
 S-CH₂-,
 φ-CH₂-O-CH₂-,
 5 tetrahydropyranyl,
 CH₃CH(-O-C₂H₅)-,
p-methoxybenzyl,
p-methoxyphenyl,
p-nitrobenzyl,
 10 (φ)₃C-,
 (CH₃)₃Si-,
 [CH₃-CH(CH₃)]₃Si- and
 φ(CH₃)₂Si-.


15 18. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 1 where the protected form of R₁ is the amino protecting group selected from the group consisting of:

(I) C₁-C₄ alkyl,
 (II) φ-CH₂-,
 20 (III) (φ)₃C-,
 (IV) R_a-CO- where R_a is (A) H-, (B) C₁-C₄ alkyl, (C) C₅-C₇ cycloalkyl, (D) (C₁-C₅ alkyl)-O-, (E) Cl₃C-CH₂-O-, (F) H₂C=CH-CH₂-O-, (G) φ-CH=CH-CH₂-O-, (H) φ-CH₂-O-, (I) *p*-methoxyphenyl-CH₂-O-, (J) *p*-nitrophenyl-CH₂-O-, (K) φ-O-, (L) CH₃-CO-CH₂-, (M) (CH₃)₃Si-O-,
 25 (V) R_b-SO₂- where R_b is: (A) (C₁ alkyl)-, (B) φ-, (C) *p*-methylphenyl- and (D) φ-CH₂-.

30 19. A process to prepare 5-hydroxymethyl substituted oxazolidinones (III) according to claim 18 where the amino protected form of R₁ is benzyloxycarbonyl.

20. A process to prepare 5-aminomethyl substituted oxazolidinone amines of formula (VII)


5



(VII)


10 where R_1 is

15

where X^1 is -H or -F;where X^2 is -H or -F;where Q^1 is:


20

a)




25

b)




30

c)




35

d)




5

f)




10

g)




15

h)




20

i)




25

j)




30

k)




35

m)



Q^1 and X^2 taken together are:



5

where Z^1 is:

- 10 a) $-CH_2-$,
 b) $-CH(R^4)-CH_2-$,
 c) $-C(O)-$, or
 d) $-CH_2CH_2CH_2-$;

where Z^2 is:

- 15 a) $-O_2S-$,
 b) $-O-$,
 c) $-N(R^7)-$,
 d) $-OS-$, or
 e) $-S-$;

where Z^3 is:

- 20 a) $-O_2S-$,
 b) $-O-$,
 c) $-OS-$, or
 d) $-S-$;

where A^1 is:

- 25 a) H- or
 b) CH_3 ;

where A^2 is:



- 30 a) H-,
 b) HO-,
 c) CH_3- ,
 d) CH_3O- ,
 e) $R^2O-CH_2-C(O)-NH-$,
 f) $R^3O-C(O)-NH-$,
 g) $(C_1-C_2)alkyl-O-C(O)-$,
 h) $HO-CH_2-$,
 35

- i) $\text{CH}_3\text{O-NH-}$,
- j) $(\text{C}_1\text{-C}_3)\text{alkyl-O}_2\text{C-}$
- k) $\text{CH}_3\text{-C(O)-}$,
- l) $\text{CH}_3\text{-C(O)-CH}_2\text{-}$,

5



10

15 A^1 and A^2 taken together are:

20

25 where R^1 is:

- a) $-\text{CHO}$,
- b) $-\text{COCH}_3$,
- c) $-\text{COCHCl}_2$,
- d) $-\text{COCHF}_2$,
- 30 e) $-\text{CO}_2\text{CH}_3$,
- f) $-\text{SO}_2\text{CH}_3$, or
- g) $-\text{COCH}_2\text{OH}$;

where R^2 is:

- a) H- ,
- 35 b) $\text{CH}_3\text{-}$,

c) phenyl-CH₂-, or

d) CH₃C(O)-;

where R³ is:

a) (C₁-C₃)alkyl-, or

5 b) phenyl-;

where R⁴ is:

a) H-, or

b) HO-;

where R⁵ is:

10 a) H-,

b) (C₁-C₃)alkyl-,

c) CH₂=CH-CH₂- or

d) CH₃-O-(CH₂)₂-;

where R⁶ is:

15 a) CH₃-C(O)-,


b) H-C(O)-,

c) Cl₂CH-C(O)-,


d) HOCH₂-C(O)-,

e) CH₃SO₂-,

20




25 g) F₂CHC(O)-,



i) H₃C-C(O)-O-CH₂-C(O)-,

30 j) H-C(O)-O-CH₂-C(O)-,




l) HC≡CH-CH₂O-CH₂-C(O)- or

35 m) phenyl-CH₂-O-CH₂-C(O)-;


where R^7 is:

- a) $R^2O-C(R^{10})(R^{11})-C(O)-$,
- b) $R^3O-C(O)-$,
- c) $R^8-C(O)-$,

5




10



15

- f) $H_3C-C(O)-(CH_2)_2-C(O)-$,
- g) R^9-SO_2- ,

20



25

- i) $HO-CH_2-C(O)-$,
- j) $R^{16}-(CH_2)_2-$,
- k) $R^{13}-C(O)-O-CH_2-C(O)-$,
- l) $(CH_3)_2N-CH_2-C(O)-NH-$,
- m) $NC-CH_2-$ or
- n) $F_2-CH-CH_2-$;

30 where R^8 is:

- a) $H-$,
- b) (C_1-C_4) alkyl,
- c) aryl $-(CH_2)_p$,
- d) ClH_2C- ,
- e) Cl_2HC- ,

35

- f) $\text{FH}_2\text{C}-$,
- g) $\text{F}_2\text{HC}-$ or
- h) $(\text{C}_3-\text{C}_6)\text{cycloalkyl}$;

where R^9 is:

- 5 a) $-\text{CH}_3$,
- b) $-\text{CH}_2\text{Cl}$,
- c) $-\text{CH}_2\text{CH}=\text{CH}_2$,
- d) aryl or
- e) $-\text{CH}_2\text{CN}$;

10 where R^{10} is H- or CH_3 -;

where R^{11} is H- or CH_3 -;

where R^{12} is:

- a) H-,
- b) $\text{CH}_3\text{O}-\text{CH}_2\text{O}-\text{CH}_2-$ or
- 15 c) HOCH_2- ;

where R^{13} is:

- a) CH_3- ,
- b) HOCH_2- ,
- c) $(\text{CH}_3)_2\text{N-phenyl}$, or
- 20 d) $(\text{CH}_3)_2\text{N}-\text{CH}_2-$;

where R^{14} is:

- a) $\text{HO}-$,
- b) $\text{CH}_3\text{O}-$,
- c) $\text{H}_2\text{N}-$,
- 25 d) $\text{CH}_3\text{O}-\text{C}(\text{O})-\text{O}-$,
- e) $\text{CH}_3-\text{C}(\text{O})-\text{O}-\text{CH}_2-\text{C}(\text{O})-\text{O}-$,
- f) $\text{phenyl}-\text{CH}_2-\text{O}-\text{CH}_2-\text{C}(\text{O})-\text{O}-$,
- g) $\text{HO}-(\text{CH}_2)_2-\text{O}-$,
- h) $\text{CH}_3\text{O}-\text{CH}_2-\text{O}-(\text{CH}_2)_2-\text{O}-$, or
- 30 i) $\text{CH}_3\text{O}-\text{CH}_2-\text{O}-$;

where R^{15} is:

- a) H- or
- b) $\text{Cl}-$;

where R^{16} is:

- 35 a) $\text{HO}-$

b) $\text{CH}_3\text{O}-$, or

c) F ;

where m is 0 or 1;

where n is 1 thru 3;

5 where p is 0 or 1;

where aryl is phenyl substituted with zero (0) or one (1) of the following:

a) $-\text{F}$,

b) $-\text{Cl}$,

c) $-\text{OCH}_3$,

10 d) $-\text{OH}$,

e) $-\text{NH}_2$,

f) $-(\text{C}_1-\text{C}_4)\text{alkyl}$,

g) $-\text{O}-\text{C}(\text{O})-\text{OCH}_3$, or

h) $-\text{NO}_2$ and protected forms thereof,


15 which comprises:

(1) contacting 5-hydroxymethyl substituted oxazolidinone alcohols of formula


(III)

(III)

20



25 where R_1 is as defined above with a sulfonylating agent selected from the group consisting of compounds of formula (V_a - V_d)



30



where n_1 is 0, 1 or 2;

where n_2 is 0 thru 4 with the provisos that:

if n_1 is 0, n_2 is 2, 3 or 4,

if n_1 is 1, n_2 is 0 or 1,


35

if n_1 is 2, n_2 is 0;

where n_3 is $5 - (n_1 + n_2)$;


where M_3 is Cl- or Br- to produce the corresponding oxazolidinone sulfonate of formula (VI_a-VI_d)

5




(VIa or VIb)

10



(VIc)

15



(VIId)

and

20 (2) contacting the oxazolidinone sulfonate (VI_a-VI_d) with ammonia at a pressure of less than about 30 psig.

21. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 20 where the sulfonylating agent (V) is selected from the group
25 consisting of 2-nitrobenzenesulfonate, 3-nitrobenzenesulfonate, 4-nitrobenzenesulfonate, 2,4-dinitrobenzenesulfonate and 2,5-dichlorobenzenesulfonate.

22. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 21 where the sulfonylating agent (V) is 3-nitrobenzenesulfonate.
30

23. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 20 where step (2) is performed at about 0 to about 20 psig.

35 24. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII)

according to claim 23 where step (2) is performed at about 0 to about 5 psig.

25. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 20 where step (2) is performed at about 60° or less.

5

26. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 20 where R₁ is phenyl substituted with one -F and one substituted amino group.

10 27. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 26 where R₁ is:

3-fluoro-4-[4-(benzyloxycarbonyl)-1-piperazinyl]phenyl or
3-fluoro-4-(4-morpholinyl)phenyl.

15 28. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 20 where step (2) is performed in the presence of an aldehyde.

29. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 28 where the aldehyde is Ar-CHO where Ar- is phenyl- optionally
20 substituted with F-, Cl-, Br-, C₁-C₅ alkyl, HO-, O₂N-, CH₃-O- or C₂H₅-O-.

30. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 28 where the aldehyde is salicylaldehyde.


25 31. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 20 where M₃ is Cl-.

32. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 20 where step (2) is performed at atmospheric pressure.


30

33. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII) according to claim 20 where the contacting is performed in the presence of water.

34. A process to prepare 5-aminomethyl substituted oxazolidinone amines (VII)
35 according to claim 20 where the sulfonylating agent (V) is




35. An oxazolidinone sulfonate of formula (VIa or VIb)



where R_1 is

10



15


where X^1 is -H or -F;

where X^2 is -H or -F;

where Q^1 is:


20

a)




25

b)




30


c)



35



Q^1 and X^2 taken together are:



5

where Z^1 is:

- 10 a) $-CH_2-$,
 b) $-CH(R^4)-CH_2-$,
 c) $-C(O)-$, or
 d) $-CH_2CH_2CH_2-$;

where Z^2 is:

- 15 a) $-O_2S-$,
 b) $-O-$,
 c) $-N(R^7)-$,
 d) $-OS-$, or
 e) $-S-$;

where Z^3 is:

- 20 a) $-O_2S-$,
 b) $-O-$,
 c) $-OS-$, or
 d) $-S-$;

where A^1 is:


- 25 a) H- or
 b) CH_3- ;

where A^2 is:


- 30 a) H-,
 b) HO-,
 c) CH_3- ,
 d) CH_3O- ,
 e) $R^2O-CH_2-C(O)-NH-$,
 f) $R^3O-C(O)-NH-$,
 g) $(C_1-C_2)_{alkyl}-O-C(O)-$,
 h) $HO-CH_2-$,
 35

- i) $\text{CH}_3\text{O-NH-}$,
 j) $(\text{C}_1\text{-C}_3)\text{alkyl-O}_2\text{C-}$
 k) $\text{CH}_3\text{-C(O)-}$,
 l) $\text{CH}_3\text{-C(O)-CH}_2\text{-}$,

5




10



A^1 and A^2 taken together are:

15



20



where R^1 is:

25

- a) $-\text{CHO}$,
 b) $-\text{COCH}_3$,
 c) $-\text{COCHCl}_2$,
 d) $-\text{COCHF}_2$,
 e) $-\text{CO}_2\text{CH}_3$,
 f) $-\text{SO}_2\text{CH}_3$, or
 g) $-\text{COCH}_2\text{OH}$;

30

where R^2 is:

- a) H- ,
 b) $\text{CH}_3\text{-}$,
 c) $\text{phenyl-CH}_2\text{-}$, or

35

d) $\text{CH}_3\text{C(O)-}$;

where R^3 is:

a) $(\text{C}_1\text{-C}_3)\text{alkyl-}$, or

b) phenyl-;

5 where R^4 is:

a) H-, or

b) HO-;

where R^5 is:

a) H-,

10 b) $(\text{C}_1\text{-C}_3)\text{alkyl-}$,

c) $\text{CH}_2 = \text{CH-CH}_2\text{-}$ or

d) $\text{CH}_3\text{-O-(CH}_2\text{)}_2\text{-}$;

where R^6 is:

a) $\text{CH}_3\text{-C(O)-}$,


15 b) H-C(O)- ,

c) $\text{Cl}_2\text{CH-C(O)-}$,

d) $\text{HOCH}_2\text{-C(O)-}$,

e) $\text{CH}_3\text{SO}_2\text{-}$,

20



g) $\text{F}_2\text{CHC(O)-}$,




25

i) $\text{H}_3\text{C-C(O)-O-CH}_2\text{-C(O)-}$,

j) $\text{H-C(O)-O-CH}_2\text{-C(O)-}$,

k)



30

l) $\text{HC=CH-CH}_2\text{O-CH}_2\text{-C(O)-}$ or


m) phenyl- $\text{CH}_2\text{-O-CH}_2\text{-C(O)-}$;

where R^7 is:


a) $\text{R}^2\text{O-C(R}^{10}\text{)(R}^{11}\text{)-C(O)-}$,

35



b) $\text{R}^3\text{O-C(O)-}$,




5






10




15



20









25




where R^8 is:

30



35



where R^9 is:

- a) $-\text{CH}_3$,
- b) $-\text{CH}_2\text{Cl}$,
- c) $-\text{CH}_2\text{CH}=\text{CH}_2$,
- 5 d) aryl or
- e) $-\text{CH}_2\text{CN}$;

where R^{10} is H- or CH_3 -;

where R^{11} is H- or CH_3 -;

where R^{12} is:

- 10 a) H-,
- b) $\text{CH}_3\text{O}-\text{CH}_2\text{O}-\text{CH}_2$ - or
- c) HOCH_2 -;

where R^{13} is:

- 15 a) CH_3 -,
- b) HOCH_2 -,
- c) $(\text{CH}_3)_2\text{N-phenyl}$, or
- d) $(\text{CH}_3)_2\text{N}-\text{CH}_2$ -;

where R^{14} is:

- 20 a) HO-,
- b) CH_3O -,
- c) H_2N -,
- d) $\text{CH}_3\text{O}-\text{C}(\text{O})-\text{O}$ -,
- e) $\text{CH}_3-\text{C}(\text{O})-\text{O}-\text{CH}_2-\text{C}(\text{O})-\text{O}$ -,
- f) $\text{phenyl}-\text{CH}_2-\text{O}-\text{CH}_2-\text{C}(\text{O})-\text{O}$ -,
- 25 g) $\text{HO}-(\text{CH}_2)_2-\text{O}$ -,
- h) $\text{CH}_3\text{O}-\text{CH}_2-\text{O}-(\text{CH}_2)_2-\text{O}$ -, or
- i) $\text{CH}_3\text{O}-\text{CH}_2-\text{O}$;

where R^{15} is:

- a) H- or
- 30 b) Cl-;

where R^{16} is:

- a) HO-
- b) CH_3O -, or
- c) F;

35 where m is 0 or 1;

where n is 1 thru 3;

where p is 0 or 1;

where aryl is phenyl substituted with zero (0) or one (1) of the following:

- a) -F,
- 5 b) -Cl,
- c) -OCH₃,
- d) -OH,
- e) -NH₂,
- f) -(C₁-C₄)alkyl,
- 10 g) -O-C(O)-OCH₃, or
- h) -NO₂ and protected forms thereof,

where n₁ is 0, 1 or 2;

where n₂ is 0 thru 4 with the provisos that:

- if n₁ is 0, n₂ is 2, 3 or 4,
 - 15 if n₁ is 1, n₂ is 0 or 1,
 - if n₁ is 2, n₂ is 0;
- where n₃ is 5 - (n₁ + n₂).

36. An oxazolidinone sulfonate (VIa or VIb) according to claim 35 which is:
- 20 3-nitrobenzenesulfonate ester (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxo)piperazinyl]-phenyl]-2-oxo-5-oxazolidinyl]methanol,
 - 2-nitrobenzenesulfonate ester (R)-[N-3-[3-fluoro-4-[N-1-(4-carbobenzoxo)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol,
 - 25 2,4-dinitrobenzenesulfonate ester (R)-[N-3-[3-fluoro-4-[N-1-(4-carbobenzoxo)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol,
 - (R)-[N-3-[3-fluoro-4-[N-1-(4-carbobenzoxo)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol 4-chlorobenzenesulfonate ester,
 - (R)-[N-3-[3-fluoro-4-[N-1-(4-carbobenzoxo)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol 2,5-dichlorobenzenesulfonate ester,
 - 30 (R)-[N-3-[3-fluoro-4-[N-1-(4-carbobenzoxo)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol 4-nitrobenzenesulfonate ester,
 - (R)-[N-3-[3-fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 3-nitrobenzenesulfonate ester,
 - (R)-[N-3-[3-fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 4-
 - 35 nitrobenzenesulfonate ester,

- (R)-[N-3-[3-fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 2-nitrobenzenesulfonate ester,
- (R)-[N-3-[3-fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 2,4-dinitrobenzenesulfonate ester,
- 5 (R)-[N-3-[3-fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 4-chlorobenzenesulfonate ester,
- (R)-[N-3-[3-fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 2,5-dichlorobenzenesulfonate ester.
- 10 37. An oxazolidinone sulfonate (VIa or VIb) according to claim 36 which is:
- 3-nitrobenzenesulfonate ester (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]-phenyl]-2-oxo-5-oxazolidinyl]methanol,
- 2-nitrobenzenesulfonate ester (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol,
- 15 2,4-dinitrobenzenesulfonate ester (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol,
- (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol 4-chlorobenzenesulfonate ester,
- (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-
- 20 oxazolidinyl]methanol 2,5-dichlorobenzenesulfonate ester,
- (R)-[N-3-[3-fluoro-4-(N-1-(4-carbobenzoxy)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methanol 4-nitrobenzenesulfonate ester,
- (R)-[N-3-[3-fluoro-4-morpholinylphenyl]-2-oxo-5-oxazolidinyl]methanol 3-nitrobenzenesulfonate ester.

25

INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/US 97/03458

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D263/24 C07D263/20 C07D413/10 C07D487/04 C07D413/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 25106 A (THE UPJOHN COMPANY) 21 September 1995 see page 12 scheme 1, pages 13, 14 see claims	1-35
A	WO 95 07271 A (THE UPJOHN COMPANY) 16 March 1995 cited in the application see page 28, chart 1, pages 12-15, example 1 see claims	1-35
A	US 4 150 029 A (DOSTERT PHILIPPE L ET AL) 17 April 1979 cited in the application see the whole document	1

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

19 June 1997

Date of mailing of the international search report

27.06.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Henry, J

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/US 97/03458

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 677 515 A (NIPPON PAINT CO LTD) 18 October 1995 see the whole document	1
P,X	WO 96 35691 A (PHARMACIA AND UPJOHN COMPANY) 14 November 1996 see page 18, line 25 - line 35; claims	1,35

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 97/ 03458

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
Claims searched completely: 36, 37
Claims searched incompletely: 1-35

see next sheet

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 97/ 03458

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

The formulation of claims 1,20,35 is so complicated because of the long lists of cascading substituents that does not comply with Art. 6 PCT prescribing that claims shall be clear and concise. For these reasons the search has been limited to the examples.

INTERNATIONAL SEARCH REPORT

Information on patent family members

In. .ional Application No

PCT/US 97/03458

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9525106 A	21-09-95	JP 8073455 A	19-03-96
		AU 2099995 A	03-10-95
		CA 2183972 A	21-09-95
		CN 1143961 A	26-02-97
		EP 0750618 A	02-01-97
WO 9507271 A	16-03-95	AU 7557094 A	27-03-95
		CA 2168560 A	16-03-95
		CN 1130379 A	04-09-96
		EP 0717738 A	26-06-96
		JP 9502436 T	11-03-97
		ZA 9405894 A	05-02-96
US 4150029 A	17-04-79	FR 2342729 A	30-09-77
		FR 2356422 A	27-01-78
		AU 509376 B	08-05-80
		AU 2268977 A	31-08-78
		BE 851893 A	29-08-77
		CA 1106380 A	04-08-81
		CH 627460 A	15-01-82
		CH 618975 A	29-08-80
		DE 2708236 A	08-09-77
		GB 1546741 A	31-05-79
		GB 1546742 A	31-05-79
		GB 1546743 A	31-05-79
		GB 1546744 A	31-05-79
		JP 1314605 C	28-04-86
		JP 52113959 A	24-09-77
		JP 60040428 B	11-09-85
		LU 76848 A	12-09-77
		NL 7702148 A	05-09-77
		SE 435928 B	29-10-84
		SE 7702013 A	02-09-77
		ZA 7701085 A	25-01-78
EP 0677515 A	18-10-95	JP 5065330 A	19-03-93
		JP 5059023 A	09-03-93
		JP 5059021 A	09-03-93
		JP 5065341 A	19-03-93
		DE 69210919 D	27-06-96

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/03458

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0677515 A		DE 69210919 T	23-01-97
		EP 0530812 A	10-03-93
		US 5324797 A	28-06-94
		US 5237021 A	17-08-93

WO 9635691 A	14-11-96	AU 5484996 A	29-11-96
